



PMI RESEARCH & DEVELOPMENT

Study ZRHM-PK-05-JP **Clinical Study Report Appendix 16.1.1** **Protocol and Protocol Amendments**

Study Title:	A single-center, open-label, randomized, controlled, crossover study to investigate the nicotine pharmacokinetic profile and safety of Tobacco Heating System 2.2 Menthol (THS 2.2 Menthol) following single use in smoking, healthy subjects compared to menthol conventional cigarettes and nicotine gum
Short Title:	Nicotine pharmacokinetic profile and safety of Tobacco Heating System 2.2 Menthol (THS 2.2 Menthol)
Study Number:	ZRHM-PK-05-JP
Product Name:	Tobacco Heating System 2.2 Menthol (THS 2.2 Menthol)
Study Initiated (first subject screened):	01 August 2013
Study Completed (last subject last visit):	16 November 2013
Principal Investigator and Affiliation:	F. Nobuoka, MD Ageo Medical Clinic 3133 Haraichi, Ageo City Saitama 362-0021, Japan
Sponsor:	Philip Morris Products S.A. PMI Research & Development Quai Jeanrenaud 5 2000 Neuchâtel, Switzerland
Sponsor Signatories:	Christelle Haziza, PhD, Manager P1 Clinical Program, Clinical Scientist Andrea Donelli, Clinical Scientist Nicola Lama, PhD, Biostatistician Patrick Picavet, MD, Medical Safety Officer
Version:	1.0
Date:	12 May 2015

This study was conducted in accordance with Good Clinical Practice.

Confidentiality Statement

This document is confidential. Disclosure of any of its contents to third parties is not permitted except by the prior written consent of Philip Morris Products S.A.



Table of Contents

16.1.1.1 STUDY PROTOCOL (ENGLISH VERSION)	3
16.1.1.2 STUDY PROTOCOL (LOCAL LANGUAGE VERSION).....	131
16.1.1.3 PROTOCOL AMENDMENTS	250
16.1.1.4 NOTES TO FILE.....	251



16.1.1.1 STUDY PROTOCOL (ENGLISH VERSION)



PMI RESEARCH & DEVELOPMENT

Clinical Study Protocol

ZRHM-PK-05-JP

Study title: A single-center, open-label, randomized, controlled, crossover study to investigate the nicotine pharmacokinetic profile and safety of Tobacco Heating System 2.2 Menthol (THS 2.2 Menthol) following single use in smoking, healthy subjects compared to menthol conventional cigarettes and nicotine gum

Product name: Tobacco Heating System 2.2 Menthol (THS 2.2 Menthol)

Short name Nicotine pharmacokinetic profile and safety of the Tobacco Heating System 2.2 Menthol (THS 2.2 Menthol)

Sponsor: Philip Morris Products S.A.
Quai Jeanrenaud 5
2000 Neuchâtel
Switzerland

Version number: Final 1.0

Date: 21 June 2013

Authors: Christelle Haziza, PhD, Manager Clinical Science
Nicola Lama, PhD, Biostatistician
Tamara Koval, MD, Medical Safety Officer
Lars Olsen, Medical Writer

Confidentiality Statement

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

SYNOPSIS

Sponsor:

Philip Morris Products S.A.

Study Title:

A single-center, open-label, randomized, controlled, crossover study to investigate the nicotine pharmacokinetic profile and safety of Tobacco Heating System 2.2 Menthol (THS 2.2 Menthol) following single use in smoking, healthy subjects compared to menthol conventional cigarettes and nicotine gum

Name of Product:

Tobacco Heating System 2.2 Menthol (THS 2.2 Menthol).

Study Number and Acronym:

ZRHM-PK-05-JP, no acronym

Primary Objective:

- To evaluate the rate and the amount of nicotine absorbed (as assessed by maximum plasma concentration [C_{\max}] and area under the concentration-time curve [AUC] from start of product use to time of last quantifiable concentration [$AUC_{(0-\text{last})}$]) from THS 2.2 Menthol relative to menthol conventional cigarettes (mCC), following single use of THS 2.2 Menthol and mCC.

Secondary Objectives:

- To determine if C_{\max} and $AUC_{(0-\text{last})}$ of plasma nicotine of the THS 2.2 Menthol are higher relative to nicotine replacement therapy gum (NRT gum) following single use of the THS 2.2 Menthol and NRT gum.
- To evaluate the difference on nicotine pharmacokinetic (PK) absorption parameters (AUC from start of product use extrapolated to time of last quantifiable concentration to infinity [$AUC_{(0-\infty)}$] and partial AUC, where t' is the subject-specific time of maximum nicotine concentration following single use of the mCC or NRT gum [$AUC_{(0-t')}$]) between the THS 2.2 Menthol and mCC, as well as the THS 2.2 Menthol and NRT gum.
- To evaluate the time to the maximum concentration (t_{\max}) of plasma nicotine for the THS 2.2 Menthol as compared to mCC and to determine if the t_{\max} for THS 2.2 Menthol is shorter as compared to NRT gum.
- To describe the terminal half-life ($t_{1/2}$) of nicotine for the THS 2.2 Menthol, mCC, and NRT gum.
- To describe the differences on urge-to-smoke over time between the THS 2.2 Menthol

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

and mCC, as well as between the THS 2.2 Menthol and NRT gum.

- To describe product evaluation in the THS 2.2 Menthol and mCC users.
- To describe the levels of carbon monoxide (CO) exposure for the THS 2.2 Menthol, as compared to mCC and NRT gum users.
- To monitor the safety during the study.

Primary Endpoints:

- Primary nicotine PK parameters (THS 2.2 Menthol vs. mCC):
 - C_{\max} .
 - $AUC_{(0-\text{last})}$.

Evaluation criterion: The study will be considered successful if the 95% confidence intervals (CI) of the THS 2.2 Menthol:mCC ratio for the primary nicotine PK parameters are estimated with a precision of $\pm 20\%$.

Secondary Endpoints:

- Primary nicotine PK parameters (THS 2.2 Menthol vs. NRT gum)
- Secondary nicotine PK parameters:
 - $AUC_{(0-\infty)}$.
 - Partial $AUC_{(0-t^*)}$.
 - t_{\max} .
 - $t_{1/2}$.
- Subjective smoking effects:
 - Urge-to-smoke questionnaire (Questionnaire of Smoking Urges brief [QSU-brief]).
 - Product evaluation questionnaire (Modified Cigarette Evaluation Questionnaire [MCEQ]).
- CO exposure biomarkers: levels of exhaled CO and carboxyhemoglobin (COHb) in blood.
- Safety variables:
 - Incidence of adverse events (AEs)/serious adverse events (SAEs), and including THS 2.2 Menthol malfunction/misuse.
 - Respiratory symptoms: cough assessment by visual analogue and Likert scales and one open question.
 - Vital signs.

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

- Spirometry.
- Electrocardiogram (ECG).
- Clinical chemistry, hematology, and urine analysis safety panel.
- Physical examination.
- Concomitant medications.

Additional Study Assessments:

- Serology for human immunodeficiency virus 1/2 and Hepatitis B and C.
- Urine pregnancy test (females only), urine cotinine test, urine drug screen.
- Alcohol breath test.
- Chest X-ray.
- Nicotine dependence to be assessed with the Fagerström Test for Nicotine Dependence revised version.
- Cytochrome P450 2A6 (CYP2A6) activity (nicotine metabolic molar ratio) in plasma.


Study Design:

This is a randomized, controlled, 2-period, 4-sequence, single-use crossover study. An incomplete block design is adopted, where each subject will receive only two of the three products ([Figure S1](#)):

- THS 2.2 Menthol.
- mCC.
- NRT gum.

Figure S1: Study Design

- Cross over with incomplete block design, 4 sequences
- 62 smokers to be randomized



Abbreviations: THS: Tobacco Heating System; mCC = menthol conventional cigarette; NRT = Nicotine Replacement Therapy

A Screening Visit will be conducted within 4 weeks prior to Admission to the investigational site (Day -29 to Day -2) (Figure S1). A demonstration of the THS 2.2 Menthol product and the NRT gum will also be done by the site study collaborator during the Screening Visit. Subjects will be admitted to the clinic on Day -1. On Day -1, as the last procedure of the eligibility assessments on that day, all subjects will undergo a product test prior to enrolment, first THS 2.2 Menthol (using of up to 3 Menthol Tobacco Sticks) and subsequently NRT gum. In female subjects, the urine pregnancy test must be negative before any product test is performed with either the THS 2.2 Menthol or the NRT gum. After all inclusion and exclusion criteria have been satisfactorily met, only subjects willing and ready to use both the THS 2.2 Menthol and the NRT gum products can be enrolled. Screening procedures do not necessarily have to be conducted on the same day.

The confinement period will then consist of 2 periods (Period 1, Period 2) with each period consisting of at least 24-hour nicotine wash-out (nicotine abstinence) and 1 day of single product use.

Period 1: Day 0: Wash-out; Day 1: single product use (THS 2.2 Menthol/mCC/NRT gum).

Period 2: Day 2: Wash-out; Day 3: single product use (THS 2.2 Menthol/mCC/NRT gum).

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

In total, 62 eligible, healthy smoking subjects will be randomized into one of the 4 sequences:

- | | | |
|-----------------------------|---|------------------------|
| Sequence 1: THS 2.2 Menthol | → | mCC (N=22) |
| Sequence 2: mCC | → | THS 2.2 Menthol (N=22) |
| Sequence 3: THS 2.2 Menthol | → | NRT gum (N=9) |
| Sequence 4: NRT gum | → | THS 2.2 Menthol (N=9) |

Subjects will be discharged (time of discharge) from the investigational site in the morning of Day 4 after performance of the Day of Discharge assessments.

From the time of discharge until Day 11: A 7-day safety follow-up will be done for the recording of spontaneously reported new AEs and SAEs, and the active follow-up of ongoing AEs/SAEs by the site.

Study Population and Main Criteria for Inclusion:

A total of 62 smoking, but healthy adult Japanese subjects, meeting the following main inclusion criteria:

- Subject is aged from 23 to 65 years (inclusive).
- Smoking, healthy subject as judged by the Investigator.
- Subject is Japanese.
- Subject smokes at least 10 commercially available mCC per day (no brand restrictions) for the last 4 weeks, based on self-reporting.
- Subject does not plan to quit smoking in the next 3 months.
- Subject is ready to accept interruptions of smoking for up to 4 days.
- Subject is ready to accept using the THS 2.2 Menthol and the NRT gum product.

Each sex and each of the smoking strata (International Organization for Standardization [ISO] nicotine levels ≤ 0.6 mg and 0.6 to 1 mg) will have a quota applied to ensure they represent at least 40% of the total study population. This will be applied to subjects randomized to sequence 1 and 2 and independently to subjects randomized to sequence 3 and 4. Subjects who do not complete the study after randomization will not be replaced.

Investigational Products, Dose, and Mode of Use:

- Test Product: THS 2.2 Menthol
- Reference Product, Dose, and Mode of Use:

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

Subject's own supply of commercially available preferred single brand mCC.

Reference Point Product (non-investigational):

Nicotine replacement therapy gum 2 mg Nicorette[®] chewed for 35 ± 5 minutes. This will be supplied by the site and reimbursed by the Sponsor or designated Clinical Research Organization (CRO).

Duration of Study:

The entire study will last 14 to 41 days, including a Screening period of up to 4 weeks prior to Admission (Day -29 to Day -2), 6 days of confinement (Day -1 to time of discharge on Day 4), and 7 days of safety follow-up (from time of discharge until Day 11).

Expected Duration of the Entire Study:

From 1st of July 2013 to 30th November 2013.

Statistical Methods:

All primary and secondary endpoints will be summarized with descriptive statistics. In addition, PK, subjective effects of smoking, and safety variables will be analyzed as follows.

Pharmacokinetics: the analysis populations for the PK endpoints will be composed of two analysis sets to allow the comparison between THS 2.2 Menthol and NRT gum separately from the comparison between THS 2.2 Menthol and mCC. Only subjects without major protocol deviations will be included in the PK analysis sets.

Nicotine PK parameters will be derived from plasma nicotine versus time data using a non-compartmental technique.

An analysis of variance (ANOVA) will be conducted on logarithmically transformed $AUC_{(0-last)}$ and C_{max} primary endpoints. The model will include terms for sequence, subjects within sequence, period, and exposure group as fixed effect factors. The results of this analysis for each of $AUC_{(0-last)}$ and C_{max} will be presented in terms of adjusted geometric least square means and 95% CIs for the THS 2.2 Menthol:mCC and THS 2.2 Menthol:NRT gum ratios. The lower bound of the 95% CI of the THS 2.2 Menthol:NRT gum ratio for C_{max} and $AUC_{(0-last)}$ will be compared with 1.00, to determine if the rate and the amount of nicotine absorbed of the THS 2.2 Menthol are higher relative to NRT gum.

The $AUC_{(0-\infty)}$, $AUC_{(0-t^*)}$, and $t_{1/2}$ will be analyzed using the same approach adopted for the primary endpoints. The one-sided Wilcoxon Signed-Rank Test ($\alpha=0.025$) will be used to test if the t_{max} in THS 2.2 Menthol is shorter than in NRT gum. The median t_{max} differences between THS 2.2 Menthol and mCC, as well as between THS 2.2 Menthol and NRT gum will be presented together with Hodges-Lehmann estimates of the 95% CI.

Subjective effects of smoking: mixed effects ANOVA using period, sequence, and product exposure as fixed effects and subjects within sequence as random effects will be adopted to

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

analyze the domain scores of the MCEQ questionnaire, for the comparison between THS 2.2 Menthol and mCC. The same model will be evaluated for the analysis of QSU-brief, including the assessment time points as repeated measurements. The results will be presented in terms of least square means and 95% CI for the THS 2.2 Menthol-mCC and THS 2.2 Menthol-NRT gum differences.

Safety: The safety population will comprise all subjects who are exposed to THS 2.2 Menthol during the study, including the THS 2.2 Menthol test at admission. Adverse event data will serve as the primary assessment of safety. All safety data will be listed and tabulated by sequence and by product use.

Sample Size:

A total of 62 subjects will be randomized. This sample size is the sum of the sample size requirements for the THS 2.2 Menthol:mCC comparison and the THS 2.2 Menthol:NRT gum comparison (as described in detail below).

The anticipated within-subject coefficients of variation for nicotine C_{\max} and $AUC_{(0-\text{last})}$ are 36% and 21%, respectively.

The sample sizes of this study assume a drop-out rate no larger than 10%.

THS 2.2 Menthol:mCC comparison

A total of 44 subjects are needed to estimate the mean C_{\max} parameter ratio between THS 2.2 Menthol and mCC with a 90% probability of obtaining a margin of error (95% CI) of at most $\pm 20\%$, assuming that THS 2.2 Menthol have a nicotine C_{\max} similar to mCC (ratio equal to 1.00). This sample size is sufficient to provide 90% probability of obtaining a margin of error of at most $\pm 20\%$ for the $AUC_{(0-\text{last})}$ ratio between THS 2.2 Menthol and mCC, assuming a similar extent of nicotine absorption for the 2 products (ratio equal to 1.00).

THS 2.2 Menthol:NRT gum comparison

A total of 18 subjects are needed to estimate the mean $AUC_{(0-\text{last})}$ parameter ratio between THS 2.2 Menthol and NRT gum with a precision allowing for the lower bound of the 95% CI exceeding 1.00, with 90% power, assuming a geometrical ratio between THS 2.2 Menthol and NRT gum of 1.28. This sample size is sufficient to provide 90% power of obtaining a lower bound of the 95% CI of C_{\max} ratio between THS 2.2 Menthol and NRT gum, exceeding 1.00, assuming a geometrical mean ratio between THS 2.2 Menthol and NRT gum of 2.00.

TABLE OF CONTENTS

SYNOPSIS.....	2
TABLE OF CONTENTS.....	9
LIST OF ABBREVIATIONS AND DEFINITIONS OF TERMS.....	13
1 ETHICS AND REGULATIONS.....	19
1.1 Institutional Review Board Approval.....	19
1.2 Ethical Conduct of the Study.....	20
1.3 Subject Information and Consent.....	20
1.4 Good Clinical Practice and Regulatory Requirements.....	21
2 INTRODUCTION.....	22
2.1 Background.....	22
2.2 Purpose of the Study.....	24
2.3 Anticipated Benefits and Risks.....	24
3 STUDY OBJECTIVES.....	26
3.1 Primary Objective.....	26
3.2 Secondary Objectives.....	26
3.3 Primary Endpoints.....	27
3.4 Secondary Endpoints.....	27
3.5 Exploratory Endpoints.....	28
4 INVESTIGATIONAL PLAN.....	29
4.1 Overall Study Design and Plan.....	29
4.2 Rationale for Study Design and Control Group(s).....	31
4.3 Study Duration.....	32
4.4 Appropriateness of Measurement.....	32
5 STUDY POPULATION.....	33
5.1 Selection of Study Population.....	33
6 INVESTIGATIONAL PRODUCTS AND REFERENCE POINT PRODUCT.....	40
6.1 Description.....	40
6.2 Use of Investigational and Reference Point Products.....	42
6.3 Method for Assigning Subjects to Study Arms.....	44
6.4 Blinding.....	44
6.5 Investigational Product Accountability and Compliance.....	45
6.6 Restrictions.....	46
7 STUDY PROCEDURES.....	50

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

7.1	Informed Consent.....	50
7.2	Smoking Cessation Advice and Debriefing.....	50
7.3	Support during Smoking Abstinence/Periods of Reduced Smoking	50
7.4	Clinical Assessments	51
7.5	Biomarker Assessment.....	54
7.6	Laboratory Assessments	56
7.7	Sample Handling, Storage, and Shipment	58
7.8	Questionnaires.....	59
8	ADVERSE EVENTS.....	62
8.1	Definitions.....	62
8.2	Assessment of Adverse Events	63
8.3	Reporting of Serious Adverse Events	65
8.4	Reporting of Other Events Critical to Safety Evaluations	66
8.5	Actions and Follow-up.....	68
9	STUDY ACTIVITIES	70
9.1	Screening Visit.....	70
9.2	Admission	72
9.3	Investigational Period	75
9.4	Day of Discharge	81
9.5	Safety Follow-up Period	83
9.6	Early Termination Procedures	83
10	QUALITY CONTROL AND QUALITY ASSURANCE.....	84
10.1	Monitoring	84
10.2	Training of Study Collaborator	85
10.3	Audits and Inspections.....	85
11	DATA MANAGEMENT ACTIVITIES.....	86
11.1	Data Capture	86
11.2	Data Handling	87
12	PLANNED STATISTICAL METHODS	89
12.1	General Considerations.....	89
12.2	Determination of Sample Size and Power Consideration.....	90
12.3	Analysis Populations.....	91
12.4	Demographics and Baseline Characteristics	92
12.5	Primary Endpoints	92
12.6	Secondary Endpoints	93

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

12.7	Exploratory Analyses.....	96
12.8	Interim Analysis.....	96
13	ADMINISTRATIVE CONSIDERATIONS.....	97
13.1	Principal Investigators and Study Administrative Structure.....	97
13.2	Subject Confidentiality	99
13.3	Access to Source Documentation	100
13.4	Record Retention	100
13.5	Clinical Study Report.....	102
13.6	Financial Disclosure.....	102
13.7	Publication and Disclosure Policy	102
13.8	Insurance	103
14	REFERENCE LIST	104

LIST OF IN-TEXT TABLES AND FIGURES

Table 1	Measured aerosol fractions for the THS Menthol Tobacco Sticks	41
Table 2	Drugs and Substances Considered Interacting with CYP2A6	47
Table 3.	Clinical Laboratory Parameters for Safety Panel.....	57
Table 4	Time Schedule – Screening Visit.....	70
Table 5.	Time Schedule – Day -1 Admission	73
Table 6.	Time Schedule – Day 0 Washout.....	75
Table 7.	Time Schedule – Day 2 Washout.....	77
Table 8.	Time Schedule – Day 1 and Day 3 Single Use.....	79
Table 9.	Time Schedule – Day 4 Discharge.....	82
Figure 1.	Study Flow Chart	31

LIST OF APPENDICES

Appendix 1	Schedule of Events.....	111
Appendix 2	Participating Laboratories	115
Appendix 3	Investigational Product and Instructions for Use.....	116
Appendix 4	THS Package Label.....	117
Appendix 5	Abnormal Laboratory Values	120

LIST OF ABBREVIATIONS AND DEFINITIONS OF TERMS

Abbreviations

ADL	Activities of daily living
AE	Adverse event
ALT	Alanine aminotransferase
ANOVA	Analysis of variance
AST	Aspartate aminotransferase
AUC	Area under the concentration-time curve
$AUC_{(0-\infty)}$	Area under the concentration-time curve from start of product use extrapolated to time of last quantifiable concentration to infinity
$AUC_{(0-last)}$	Area under the concentration-time curve from start of product use to time of last quantifiable concentration
$AUC_{(0-t')}$	Partial AUC, where t' is the subject-specific time of maximum nicotine concentration following single use of mentholated conventional cigarettes or NRT gum.
BMI	Body mass index
CC	Conventional cigarette(s)
CD	Compact disc
CI	Confidence Interval
C_{last}	Last quantifiable concentration
C_{max}	Maximum plasma concentration
CO	Carbon monoxide

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

COHb	Carboxyhemoglobin
CRA	Clinical Research Associate
CRO	Contract Research Organization
CSR	Clinical Study Report
CTCAE	Common Terminology Criteria for Adverse Events and Common Toxicity Criteria
CTMS	Clinical Trial Management System
CV	Coefficient of variation
CYP2A6	Cytochrome P450 2A6
DMP	Data Management Plan
ECG	Electrocardiogram
eCRF	Electronic Case Report Form
EOS	End of Study Visit (referring to each subject's individual last study visit)
ePRO	Electronic patient reported outcome
FDA	Food and Drug Administration
FEV ₁	Forced expiratory volume in 1 second
FTND	Fagerström Test for Nicotine Dependence (revised version)
FVC	Forced vital capacity
GCP	Good Clinical Practice
HIV	Human immunodeficiency virus
HPHCs	Harmful and potentially harmful constituents

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

IB	Investigator's Brochure
ICF	Informed consent form
ICH	International Conference on Harmonization
IP	Investigational Product
IRB	Institutional Review Board
ISO	International Organization for Standardization
IV	Intravenous
LLN	Lower limit of the normal range
LLOQ	Lower limit of quantification
mCC	Menthol conventional cigarette(s)
MCEQ	Modified Cigarette Evaluation Questionnaire
MedDRA	Medical Dictionary for Regulatory Activities
MRTP	Modified risk tobacco product
n	Number of subjects
NRT gum	Nicotine replacement therapy gum
PK	Pharmacokinetic(s)
PMI	Philip Morris International
QC	Quality control
QSU-brief	Questionnaire of Smoking Urges brief
SA	Smoking abstinence

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

SAE	Serious adverse event
SAP	Statistical Analysis Plan
SHM	Sample handling manual
SOP	Standard Operating Procedure
T ₀	Time point of first product use during study day
t _{1/2}	Terminal half-life
THS 2.2 Menthol	Tobacco Heating System 2.2 Menthol
t _{max}	Time to the maximum concentration
UBC	United BioSource Corporation
ULN	Upper limit of the normal range
ULOQ	Upper limit of quantification
VAS	Visual Analogue Scale
WBC	White blood cell (count)
WHO	World Health Organization
λ _z	Terminal elimination rate constant

Explanations of Terms

The following special terms are used in this protocol:

CC and mCC	The term ‘conventional cigarette’ refers to manufactured and commercially available cigarettes and excludes hand-rolled cigarettes, cigars, pipes, bidis, and other nicotine-containing products. Menthol CC are designated “mCC”.
Charger	The function of the Charger (Model 4) is to recharge the Holder after use. It contains a battery with sufficient capacity to recharge the Holder approximately 20 times. It is a convenient size to carry around, and can itself be recharged from a mains power source.
Day of Discharge	Day 4.
End of Study	End of Study is defined as the last day of the 7-day safety follow-up subsequent to discharge from the unit.
Enrolment	On Day -1 for eligible subjects after all applicable inclusion and exclusion criteria have been satisfactorily assessed and the subject is willing and ready to use both the Tobacco Heating System THS 2.2 Menthol and nicotine replacement therapy gum (NRT gum). The test of both THS 2.2 Menthol and NRT gum are the last assessments prior to enrolment.
Investigator	Principal Investigator or sub-Investigator
First product use time point	Start of product use for THS 2.2 Menthol is defined as the time of the first puff. The start time for mCC corresponds to the lighting of the mCC, and the start time of the NRT gum product is the time of NRT gum intake.
Randomization	Assignment to product on Day 0 utilizing an Interactive Web and Voice Response System.
Safety follow-up	After the time of discharge, a 7-day safety follow-up will be done for the recording of spontaneously reported new AEs/SAEs and the active follow-up of ongoing adverse events (AEs)/serious adverse events (SAEs) by the site. In general any AE will be followed up until resolved, stabilized i.e., no worsening of the event or a plausible explanation for the event has been found.

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

Screening failure	Subjects who do not meet the entry criteria from informed consent form (ICF) signature to the time of enrolment will be considered a screening failure and will be replaced by other subjects.
THS Menthol Tobacco Stick (Menthol Tobacco Stick)	The Menthol Tobacco Stick (product code C3 menthol) contains tobacco which, when heated, generates an aerosol. It is custom-designed to be used with the Holder.
Tobacco Stick Holder (Holder)	The function of the Holder (Model 4.2) is to heat the Menthol Tobacco Stick, delivering an aerosol to the user. The electrical heating is powered from an internal battery which delivers power for about 6 minutes (allowing complete use of a single Menthol Tobacco Stick)

1 ETHICS AND REGULATIONS

1.1 Institutional Review Board Approval

Prior to the start of the study, the clinical study protocol, together with its associated documents (informed consent form [ICF] which includes both subject information sheet and informed consent, subject recruitment procedures [e.g., advertisements], written information to be provided to the subjects, Investigator's Brochure [IB], available safety information, the Principal Investigator's curriculum vitae and/or other evidence of qualifications, the list of sub-Investigators and any other documents requested by the Institutional Review Board [IRB]), will be submitted for review and approval to the relevant IRB. The IRB shall be appropriately constituted and perform its functions in accordance with the International Conference on Harmonisation (ICH) Tripartite Guidance for Good Clinical Practice (GCP), Ministerial Ordinance on GCP for Drugs, and local requirements, as applicable.

In accordance with GCP, a written confirmation of the IRB approval should be provided to the Sponsor. This should identify the study (Principal Investigator's name, study number, and title) and the documents that have been approved by the IRB, with dates and version numbers, as well as the date of approval. The composition of the IRB, including the name and occupation of the chairperson, should be supplied to the Sponsor together with a GCP compliance statement.

The written approval from the IRB will be filed in the Investigator file, and another original copy will be filed in the Study Master File at the Sponsor or designated organization. The study must not start before the Sponsor has obtained written confirmation of favorable opinion/approval from the concerned IRB.

Any substantial change or addition to this protocol will require a written protocol amendment that must be approved by the Sponsor and the Principal Investigator. All amendments will be submitted to the IRB, and substantial amendments will only be implemented after approval by the IRB.

These requirements for approval should in no way prevent any action from being taken by the Principal Investigator or sub-Investigator (Investigator) or by the Sponsor in order to eliminate immediate hazards to the subjects. If such a change to the protocol is felt to be necessary by the Investigator, and is implemented for safety reasons, the Sponsor and the IRB should be informed immediately.

Relevant safety information will be submitted to the IRB during the course of the study in accordance with national regulations and requirements.

1.2 Ethical Conduct of the Study

The study will be performed in accordance with ethical principles that have their origin in the [Declaration of Helsinki \(2008\)](#) and are consistent with ICH/GCP, Ethical Guidelines for Clinical Studies (Ministry of Health, Labour and Welfare, 2003 (as last amended on July 31, 2008)), Ministerial Ordinance on Good Clinical Practice for Drugs ([Ministry of Health, Labour and Welfare, 1997](#) (as last amended by the Ordinance of Ministry of Health, Labour and Welfare No. 161 of December 28, 2012)), and applicable regulatory principles.

The Principal Investigator agrees to conduct the clinical study in compliance with the protocol agreed upon with the Sponsor and approved by the IRB. The Principal Investigator and the Sponsor must sign the protocol (and protocol amendments, if applicable) to confirm this agreement. A copy of the [Declaration of Helsinki \(2008\)](#) should be located in the Investigator's Study File.

1.3 Subject Information and Consent

1.3.1 Study Consent/Subject Information Sheet

Before or at the Screening Visit, the Investigator will ensure each subject is given full and adequate oral and written information about the nature, purpose, possible risks and benefits of the study, and the Investigator will answer all questions the subject might have to his/her full satisfaction. The subject will have sufficient time for consideration of his/her participation in the study and will be notified that he/she is free to discontinue his/her participation at any time. Once the subject has received all necessary information, and if he/she agrees to participate, this will be documented in the ICF by the date and signature of both the subject and the person who conducted the informed consent discussion. No study-specific procedures will be performed before the ICF sheet has been signed.

The original, dated and signed ICF(s) must be kept in the Investigator file at the site, and a copy must be given to the subject.

The subject will be informed that additional data analyses not mentioned in the protocol or the statistical analysis plan might be performed with the collected data at a later time. If any additional analyses will be performed, they will fully be covered by data confidentiality, as for the main analyses described in this protocol.

1.3.2 Amendment to the Informed Consent Form/Subject Information Sheet

If a protocol amendment is required, or if new information regarding the risk profile of the Investigational Product (IP) becomes available, an amendment may be required to the ICF. If revision of the ICF is necessary, the Principal Investigator will, with the support of the Sponsor, ensure that the documents have been reviewed and approved by a relevant IRB before subjects are required to re-sign the ICF.

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

1.4 Good Clinical Practice and Regulatory Requirements

The procedures set out in this clinical study protocol pertaining to the conduct, evaluation, and documentation of this study are designed to ensure that the Sponsor, its authorized representative, and the Principal Investigator abide by the principles of the ICH guidelines on GCP. These guidelines apply specifically to pharmaceutical development, but nevertheless provide a robust and ethical framework for conducting clinical studies of tobacco products. In addition, the Investigator will carry out the clinical study in accordance with GCP and Ethical Guidelines for Clinical Studies.

2 INTRODUCTION

2.1 Background

2.1.1 Smoking-Related Diseases and Harm Reduction Strategy

Cigarette smoking causes pulmonary and cardiovascular diseases and other serious diseases in smokers (U.S. Department of Health and Human Services, 2010). The effects of smoking and smoking cessation on mortality from cardiovascular disease among the Japanese population were investigated in cohort studies in Japan. These studies confirmed the association between smoking and mortality from cardiovascular disease and highlighted the importance of smoking cessation at any age to prevent cardiovascular disease in the Japanese population ([Honjo et al. 2010a](#), [Iso et al. 2005](#)). In 2000 in Japan, the proportion of adult deaths due to smoking ranges from 20.0% to 24.9% in males and from 10.0% to 14.6% in females. In 2007, Japan became part of the top 5 conventional cigarette (CC)-consuming countries with a consumption of 234 billion annually in 2009 (World Lung Foundation, 2012). There is no safe cigarette and the best way for smokers to reduce the adverse health consequences of smoking is to quit. Despite the risks which are attributable to smoking, some smokers cannot refrain from smoking or decide to continue smoking. To those smokers who are not able or not willing to quit, Philip Morris International (PMI) is developing alternative approaches by developing products with the potential to reduce the risks of tobacco-related diseases. These products are now referred to by the US Food and Drug Administration (FDA) as modified risk tobacco products (MRTPs) ([FDA, 2012c](#)).

The challenge in developing and commercializing MRTPs is two-fold, i.e. developing tobacco products that are shown to reduce risk and that are acceptable to smokers as substitutes for CCs. PMI is developing candidate MRTP that provide an inhalation experience without combustion. The novel approach to achieve this is by heating tobacco at significantly lower temperatures than required for CC.

PMI's approach to scientifically assessing the risk-reduction potential of its candidate MRTPs is described in the reference document ([PMI White Paper Docket](#)). Smoking cessation is the only intervention proven to reduce the risk of smoking-related diseases in smokers. Accordingly, PMI utilizes smoking cessation/smoking abstinence (SA) as the benchmark for assessing the risk reduction potential of its candidate MRTPs. The Institute of Medicine observed that cessation is the "gold standard" for assessing risk reduction, and that "the closer risks and exposures from the MRTP are to cessation products, the more confident a regulator can be of achieving a net public health benefit" ([Institute of Medicine, 2012](#)). PMI has already conducted studies and plans to conduct further clinical studies which observe measurable changes in blood chemistry, risk factors and health effects in smokers who switch to a candidate MRTP, comparing the changes with those observed in both smokers who continue smoking CC and smokers who stop using tobacco products. Longer-

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

term data from adults who continue to use the candidate MRTP can further substantiate reductions in individual risk in smokers and population harm.

2.1.2 Description of the Product and Scientific Findings

Thousands of chemicals - “smoke constituents” - are formed when tobacco is burned or combusted. More than 5,300 smoke constituents have been identified ([Rodgman and Perfetti, 2009](#)), and more than 100 of them have been categorized as harmful and potentially harmful constituents (HPHC) ([FDA, 2011](#)). There is no convincing evidence that selective reduction of smoke constituents can reduce tobacco related diseases ([Hatsukami et al., 2007](#)). Philip Morris International’s focus has been the development of products that do not combust tobacco but which replicate the “smoking experience” traditionally obtained with CC, as much as possible. Our approach limits pyrolysis and combustion, by heating tobacco at significantly lower temperatures than CC. Philip Morris International believes that such products present the best opportunity for reducing harm because they produce vastly lower levels of harmful smoke constituents and are more likely to be accepted by smokers as substitutes for CCs. Important to this effort has been providing nicotine in a way that closely parallels CC.

The product developed by PMI, and to be assessed in this study, is the Tobacco Heating System 2.2 Menthol (THS 2.2 Menthol). With this product, the heating of the tobacco is maintained below 400°C, a temperature much lower than what is observed for CC, which can reach 900°C. The THS 2.2 Menthol is composed of the ‘THS Tobacco Stick Holder’, dedicated special Menthol Tobacco Sticks made of conventional tobacco, a Charger, and different accessories. The energy of the THS Tobacco Stick Holder is sufficient to maintain approximately a 6 minute session. Unlike mCC, the Menthol Tobacco Sticks do not burn down during their consumption and their lengths remain constant after use.

The non-clinical assessment of earlier development of THS 2.2 Menthol described in the Investigator’s Brochure ([PMI, 2013a](#)) supports the initiation of the clinical studies. No new or increased toxicological hazard in the product’s aerosol was detected compared with CC smoke. The aerosol was chemically analyzed confirming that none of the determined HPHCs in the THS 2.2 Menthol were increased compared to the conventional cigarettes. The biological activity was tested in a number of *in vitro* assays to assess the cytotoxicity and the genotoxicity of the aerosol fractions total particulate matter (TPM) and gas vapor phase (GVP). In vitro and in vivo results corroborated the concept that absence of combustion when consuming tobacco substantially lowers toxic effects seen in these biological models. Further details on the clinical data are provided in the Investigators’ Brochure ([PMI, 2013a](#)).

Several clinical studies have been conducted on THS 1.0 and THS 1.0 Menthol, an earlier development version of THS 2.2 Menthol, in Europe, Asia, Africa and the United States. All studies showed reductions in exposure to the majority of measured HPHCs from both aerosol fractions, TPM and GVP, in subjects who used the THS 1.0 as compared to subjects

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

continuing smoking CC, both, in controlled and ambulatory conditions. No clinical studies were conducted with THS 2.2 Menthol.

The previous version of THS 2.2 Menthol, namely THS 2.1 (non-menthol version) was tested in two exploratory clinical studies to measure the nicotine plasma kinetic profile (PK) (www.clinicaltrials.gov identifier: NCT01780688) and to assess the reduction of exposure to HPHCs (www.clinicaltrials.gov identifier: NCT01780714) when switching from CC to THS 2.1. The observed nicotine plasma PK profile for THS 2.1 was similar to CC as well, there were significant reductions in the exposure to the majority of selected HPHCs. Clinical studies conducted so far revealed no safety concern for either of the previous version of THS 2.2 Menthol tested. Further details on the clinical data are provided in the Investigators' Brochures ([PMI, 2013a](#)).

2.2 Purpose of the Study

The purpose of this clinical study is to compare the profile of nicotine uptake (rate and extent of nicotine absorbed) after single use of switching from menthol conventional cigarettes (mCC) to THS 2.2 Menthol and mCC in smoking Japanese healthy smokers. THS 2.2 Menthol will also be compared with the nicotine replacement therapy gum (NRT gum), used as a reference point.

2.3 Anticipated Benefits and Risks

2.3.1 Anticipated Benefits

In Japan, research conducted by the Ministry of Health, Labour and Welfare has shown that 35.9% of male and 43.6% of female respondents over 20 years of age expressed the wish to quit smoking ([International Tobacco Online, 2012](#)). In a large follow-up study of middle aged Japanese smokers, the predictors of smoking cessation were age, job, smoking habit, physical activity, health checkup participation, and health status ([Honjo et al. 2010b](#)). Advice on health risks associated with smoking and smoking cessation advice will be provided at Screening, at Admission, and at the Day of Discharge. The advice will follow the recommendations of the World Health Organization (WHO) ([WHO, 2002](#)) -“Evidence based Recommendations on the Treatment of Tobacco Dependence”. Subjects who are motivated to quit smoking during the study will be given the opportunity to continue their smoking cessation attempt and will be referred to appropriate stop smoking services for continuing support and counseling at a higher level. Subjects who participate in this study will also benefit from repeated, detailed health check-ups, which may help to uncover undiagnosed medical conditions.

2.3.2 Anticipated Foreseeable Risks due to Study Procedures

- Risks related to blood sampling, e.g., excessive bleeding, fainting, hematoma, paresthesia, or infection.
- Risks related to chest X-rays, e.g., a small increase of risk to develop cancer later in life.
- Risks related to drug application as part of testing procedures (i.e. spirometry with short-acting bronchodilator at Screening) per study protocol and scientifically accepted standards.

2.3.3 Anticipated Foreseeable Risks due to Investigational Product (THS 2.2 Menthol/mCC) or Reference Point Product

- Change in smoking habits due to study requirements and related concomitant symptoms, (e.g., craving, withdrawal symptoms).

All risks related to study procedures, IP, or support for SA will be explained in detail to the subjects. Mitigation will include, but will not be limited to:

- Close monitoring and medical evaluation of potential safety signals throughout the study and follow-up.
- Using accepted research and scientific standards, (e.g., blood samples not to exceed local blood donation standards).
- Management and follow-up of adverse events (AEs)/serious adverse events (SAEs).
- Risks specific to the use of any NRT gum, as per the relevant summary of product characteristics.

2.3.4 Unforeseeable Risks

As with any new IP, and the reference point product, there may be unforeseeable risks and hazards that could occur. The possibility of such will be explained at Screening and at Admission. Mitigation will include close monitoring and medical supervision to detect any unforeseeable risk or safety signals at the earliest possibility.

3 STUDY OBJECTIVES

3.1 Primary Objective

The primary objective of this study is:

- To evaluate the rate and the amount of nicotine absorbed (as assessed by maximum plasma concentration [C_{\max}] and area under the concentration-time curve [AUC] from start of product use to time of last quantifiable concentration [$AUC_{(0-\text{last})}$]) from THS 2.2 Menthol relative to mCC, following single use of THS 2.2 Menthol and mCC.

3.2 Secondary Objectives

The secondary objectives of this study are:

- To determine if C_{\max} and $AUC_{(0-\text{last})}$ of plasma nicotine of the THS 2.2 Menthol are higher relative to NRT gum following single use of the THS 2.2 Menthol and NRT gum.
- To evaluate the difference on nicotine pharmacokinetic (PK) absorption parameters (AUC from start of product use extrapolated to time of last quantifiable concentration to infinity [$AUC_{(0-\infty)}$] and partial AUC, where t' is the subject-specific time of maximum nicotine concentration following single use of mCC or NRT gum [$AUC_{(0-t')}$]) between the THS 2.2 Menthol and mCC, as well as the THS 2.2 Menthol and NRT gum.
- To evaluate the time to the maximum concentration (t_{\max}) of plasma nicotine for the THS 2.2 Menthol as compared to mCC and to determine if the t_{\max} for THS 2.2 Menthol is shorter as compared to NRT gum.
- To describe the terminal half-life ($t_{1/2}$) of nicotine for the THS 2.2 Menthol, mCC, and NRT gum.
- To describe the differences on urge-to-smoke over time between the THS 2.2 Menthol and mCC, as well as between the THS 2.2 Menthol and NRT gum.
- To describe product evaluation in the THS 2.2 Menthol and mCC users.
- To describe the levels of carbon monoxide (CO) exposure for the THS 2.2 Menthol, as compared to mCC and NRT gum users.
- To monitor the safety during the study.

3.3 Primary Endpoints

- Primary nicotine PK parameters (THS 2.2 Menthol vs. mCC):
 - C_{\max} .
 - $AUC_{(0-\text{last})}$.

Evaluation criterion:

The study will be considered successful if the 95% confidence intervals (CI) of the THS 2.2 Menthol:mCC ratio for the primary nicotine PK parameters are estimated with a precision of $\pm 20\%$.

3.4 Secondary Endpoints

- Primary nicotine PK parameters (THS 2.2 Menthol vs. NRT gum)
- Secondary nicotine PK parameters
 - $AUC_{(0-\infty)}$.
 - Partial $AUC_{(0-t^*)}$.
 - t_{\max} .
 - $t_{1/2}$.
- Subjective smoking effects:
 - Urge-to-smoke questionnaire (Questionnaire of Smoking Urges brief [QSU-brief]).
 - Product evaluation questionnaire (Modified Cigarette Evaluation Questionnaire [MCEQ]).
- CO exposure biomarkers: levels of exhaled CO and carboxyhemoglobin (COHb) in blood.
- Safety variables:
 - Incidence of AEs/SAEs and device events, including THS 2.2 Menthol malfunction/misuse.
 - Respiratory symptoms: cough assessment by Visual Analogue Scale (VAS) and Likert scales and one open question.
 - Vital signs.
 - Spirometry.
 - Electrocardiogram (ECG).
 - Clinical chemistry, hematology, and urine analysis safety panel.
 - Physical examination.
 - Concomitant medications.

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

Additional Study Assessments:

- Serology for human immunodeficiency virus 1/2 and Hepatitis B and C.
- Urine pregnancy test (females only), urine cotinine test, urine drug screen.
- Alcohol breath test.
- Chest X-ray.
- Nicotine dependence to be assessed with the Fagerström Test for Nicotine Dependence (FTND) revised version, FTND.
- Cytochrome P450 2A6 (CYP2A6) activity (nicotine metabolic molar ratio) in plasma.

3.5 Exploratory Endpoints

There are no exploratory analyses planned.

4 INVESTIGATIONAL PLAN

4.1 Overall Study Design and Plan

This is a randomized, controlled, 2-period, 4-sequence, single use crossover study where each subject will receive two of the three products:

- THS 2.2 Menthol.
- mCC.
- NRT gum.

A Screening Visit will be conducted within 4 weeks prior to Admission to the investigational site (Day -29 to Day -2) (Figure 1). A demonstration of the THS 2.2 Menthol product and the NRT gum will also be done by the site staff during the Screening Visit. Subjects will be admitted to the clinic on Day -1. On Day -1, as the last procedure of the eligibility assessments on that day, all subjects will undergo a product test prior to enrolment: first THS 2.2 Menthol and subsequently NRT gum (using up to 3 Menthol Tobacco Sticks). In female subjects, the urine pregnancy test must be negative before any product test is performed with either the THS 2.2 Menthol or the NRT gum. After all inclusion and exclusion criteria have been satisfactorily met, only subjects willing and ready to use both the THS 2.2 Menthol and the NRT gum products can be enrolled. Screening procedures do not necessarily have to be conducted on the same day.

The confinement period will then consist of 2 periods (Period 1, Period 2) with each period consisting of at least 24-hour nicotine wash-out (nicotine abstinence) and 1 day of single product use.

Period 1: Day 0: Wash-out; Day 1: single product use (THS 2.2 Menthol/mCC/NRT gum).

Period 2: Day 2: Wash-out; Day 3: single product use (THS 2.2 Menthol/mCC/NRT gum).

In total, 62 eligible, smoking subjects will be randomized into 1 of the 4 sequences:

Sequence 1: THS 2.2 Menthol	→	mCC (N=22).
Sequence 2: mCC	→	THS 2.2 Menthol (N=22).
Sequence 3: THS 2.2 Menthol	→	NRT gum (N=9).
Sequence 4: NRT gum	→	THS 2.2 Menthol (N=9).

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

This procedure will lead to an incomplete block design with every subject being exposed to 2 of the 3 study products, as the comparison between NRT gum and mCC will not be considered.


mCC vs. THS 2.2 Menthol → (N=22 in both sequence 1 and 2)
THS 2.2 Menthol vs. NRT gum → (N=9 in both sequence 3 and 4)

Each sex and each of the smoking strata (International Organization for Standardization [ISO] nicotine levels ≤ 0.6 mg and 0.6 to 1 mg) will have a quota applied to ensure they represent at least 40% of the total study population. This will be applied to subjects randomized to sequence 1 and 2 and independently to subjects randomized to sequence 3 and 4.

Subjects will be discharged (time of discharge) from the investigational site in the morning of Day 4 after all examinations of the Day of Discharge have been conducted. After the time of discharge, a 7-day safety follow-up will be started for the recording of spontaneously reported new AEs/SAEs and the active follow-up of ongoing AEs/SAEs by the site. Any AE will, in general, be followed up until resolved, stabilized (i.e., no worsening of the event), or until a plausible explanation for the event has been found.

Figure 1. Study Flow Chart

- Cross over with incomplete block design, 4 sequences
- 62 smokers to be randomized



Abbreviations: THS: Tobacco Heating System; mCC = menthol conventional cigarette; NRT = Nicotine Replacement Therapy

4.2 Rationale for Study Design and Control Group(s)

The minimum age of 23 years age in the inclusion criteria was selected based on:

- The legal age of smoking in Japan is 20 years.
- To account for the 3 years of smoking history.

In this study, mCCs (up to 1.0 mg nicotine ISO yield/mCC) will be used as a reference product and a market-approved pharmaceutical NRT gum 2 mg (Nicorette®) will be used as non-investigational reference point. The NRT gum has been selected as the reference point, because of its availability over the market. The NRT gum will serve as a reference point for comparison with THS 2.2 Menthol for the following endpoints:

- Nicotine PK parameters.
- Urge-to-smoke.
- Safety profile.

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

The nicotine wash-out period was set to at least 24 hours ($>5 \times$ elimination $t_{1/2}$) as the elimination $t_{1/2}$ of nicotine in blood is around 2 hours in Caucasian smokers (Benowitz et al, 2009).

The use of estrogen contraceptive is known to accelerate nicotine clearance by 20% to 30% in women as compared to women who do not take such contraceptives (Benowitz et al, 2006). Therefore, for the purpose of this study, it is not allowed to use hormonal contraception containing estrogens. This also applies to hormone replacement therapy.

In this study, THS 2.2 Menthol will be tested against mCC and NRT gum will be used as a reference point product.

In this study, smokers of mCC will be assessed because the menthol brands play a significant role within the Japanese market, with 20% of the overall market share in 2008 (Connolly et al., 2011).

Furthermore, CYP2A6 activity will be assessed in this study. Asians on average metabolize nicotine more slowly than Caucasians do, at least in part due to a high prevalence of the CYP2A6 alleles associated with reduced or absent enzyme activity (Matta et al. 2006).

4.3 Study Duration

The entire study per subject will last 14 to 41 days, including a Screening period of up to 4 weeks prior to Admission (Day -29 to Day -2), and 6 days of confinement (Day -1 to morning of Day 4), and 7 days of safety follow-up (from time of discharge until Day 11).. In the morning of Day 4, the Day of Discharge examinations will be conducted. After the time of discharge, subjects will then enter a 7-day safety follow-up (until Day 11) for the recording of spontaneously reported new AEs/SAEs and the active follow-up of ongoing AEs/SAEs.

4.4 Appropriateness of Measurement

All laboratory measures utilized for this study are validated and are appropriate for the study assessments.

All used questionnaires, except the cough questionnaire, are either available as a validated questionnaire in the local language or will be forward-translated and back-translated with subsequent independent verification.

5 STUDY POPULATION

5.1 Selection of Study Population

Sixty-two Japanese female or male smoking healthy adult subjects, who smoke at least 10 mCC per day will be randomized into this study. The maximum number of mCC is not limited. Subjects must have a smoking history of at least 3 years of consecutive smoking prior to Screening. The smoking status of the subjects will be verified based on a urine cotinine test (cotinine ≥ 200 ng/mL).

5.1.1 Inclusion Criteria

Inclusion Criteria	Rationale	Screening	Day of Admission (Day -1)
1. Subject has signed the ICF and is able to understand the information provided in the Subject Information Sheet and ICF.	Administrative	X	
2. Subject is aged from 23 to 65 years (inclusive).	Safety	X	
3. Subject is Japanese.	Effect	X	
4. Smoking, healthy subject as judged by the Investigator based on all available assessments in the Screening period/day of Admission (e.g., safety laboratory, spirometry* [forced expiratory volume in 1 second {FEV ₁ }/forced vital capacity {FVC} >0.7 at post-bronchodilator basal spirometry, post-bronchodilator FEV ₁ >80% predicted value, and post-bronchodilator FVC >0.8], vital signs, physical examination, ECG, chest X-ray and medical history).	Safety	X	X

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

Inclusion Criteria	Rationale	Screening	Day of Admission (Day -1)
5. Subject smokes at least 10 commercially available menthol CCs per day (no brand restrictions) with a maximum yield of 1 mg nicotine ISO/mCC, as labelled on the cigarette package, for the last 4 weeks, based on self-reporting. Furthermore, the subject has been smoking for at least the last 3 consecutive years. The smoking status will be verified based on a urinary cotinine test (cotinine ≥ 200 ng/mL).	Effect	X	X
6. The subject does not plan to quit smoking in the next 3 months.	Safety	X	
7. The subject is ready to accept interruptions of smoking for up to 4 days.	Safety	X	X
8. The subject is ready to accept using both the THS 2.2 Menthol and NRT gum products.	Effect		X

5.1.2 Exclusion Criteria

Subjects who meet any of the following exclusion criteria must not be enrolled into the study:

Exclusion Criteria	Rationale	Screening	Day of Admission (Day -1)
1. As per Investigator judgment, the subject cannot participate in the study for any reason (e.g., medical, psychiatric, and/or social reason).	Safety	X	X

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

Exclusion Criteria	Rationale	Screening	Day of Admission (Day -1)
2. A subject who is legally incompetent, physically or mentally incapable of giving consent (e.g., emergency situation, under guardianship, , prisoners or subjects who are involuntarily incarcerated).	Administrative	X	
3. The subject has medical condition requiring smoking cessation, or clinically relevant diseases (including but not limited to gastrointestinal, renal, hepatic, neurological, hematological, endocrine, oncological, urological, immunological, pulmonary and cardiovascular disease or any other medical condition [including but not limited to clinically relevant abnormal laboratory parameters]) in the judgment of the Investigator.	Safety	X	X
4. The subject has a body mass index (BMI) <18.5 or ≥ 32.0 kg/m ² .	Safety	X	X
5. As per Investigator judgment, the subject has medical conditions which require or will require in the course of the study, a medical intervention (e.g., start of treatment, surgery, hospitalization) which may interfere with the study participation and/or study results.	Effect	X	X

Exclusion Criteria	Rationale	Screening	Day of Admission (Day -1)
6. The subject has used nicotine-containing products other than commercially available mCC (either tobacco-based products or nicotine-replacement therapy) as well as electronic cigarettes and similar devices, within 4 weeks prior to assessment.	Effect	X	X
7. The subject has received medication (prescription or over-the-counter) within 14 days or within 5 half-lives of the drug prior to the Admission Day (Day -1; whichever is longer) that has an impact on CYP2A6 activity.	Effect		X
8. In case the subject received any medication (prescribed or over-the-counter) within 14 days prior to Screening or prior to the Admission Day (Day -1) it will be decided at the discretion of the Investigator if these can potentially interfere with the study objectives and subject's safety.	Effect	X	X
9. The subject has a positive alcohol test and/or the subject has a history of alcohol abuse that could interfere with subject's participation in study.	Administrative	X	X
10. The subject has a positive urine drug test.	Administrative	X	X
11. Positive serology test for human immunodeficiency virus (HIV) 1/2, Hepatitis B or Hepatitis C.	Safety	X	

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

Exclusion Criteria	Rationale	Screening	Day of Admission (Day -1)
12. Donation or receipt of whole blood or blood products within 3 months prior to Admission.	Safety	X	X
13. The subject is a current or former employee of the tobacco industry or of their first-degree relatives (parent, sibling, child).	Administrative	X	
14. The subject is an employee of the investigational site or any other parties involved in the study or of their first-degree relatives (parent, sibling, child).	Administrative	X	
15. The subject has participated in a clinical study within 3 months prior to the Screening Visit.	Safety	X	
16. The subject has previously participated in the same study at a different time (i.e., each subject can be included in the study population only once).	Administrative	X	
17. For women only: Subject is pregnant (does not have negative pregnancy tests at Screening and at Admission) or is breast feeding.	Safety	X	X
18. For women only: Subject does not agree to use an acceptable method of effective contraception.*	Safety	X	X

* Intrauterine device, intrauterine system, barrier methods of contraception (condoms, occlusive caps) with spermicidal foam/gel/film/suppository, hormonal contraception

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

containing progesterone only, vasectomized partner(s) or true abstinence (periodic abstinence and withdrawal are not effective methods) from Screening until the end of the safety follow-up period. Hormonal contraception with estrogen containing products is NOT allowed in this study.

5.1.3 Removal of Subjects from the Study

Subjects will be informed that they are free to withdraw from the study at any time. Subjects should be questioned for the reason of premature withdrawal, although they are not obliged to disclose it. This needs to be fully documented in the Source Document and electronic Case Report Form (eCRF).

When a subject withdraws or is removed from the study, the whole examination procedure planned on Day 4 must be performed as soon as possible after the time of withdrawal unless the subject has withdrawn their informed consent to do so. After the time of withdrawal, the subject will enter into the 7-day period of safety follow-up. Subjects withdrawn or removed from the study cannot re-enter the study.

Subjects must be withdrawn from the study for any of the following reasons:

- Withdrawal of informed consent.
- Any AE or condition (including clinically relevant changes in a laboratory parameter), at the discretion of the Investigator.
- Positive pregnancy testing (any invasive procedures, including the drawing of blood MUST NOT be performed after diagnosis of pregnancy, see [Section 8.5.1](#)).
- The Sponsor or Principal Investigator terminates the study. If the Sponsor or the Principal Investigator decide to prematurely terminate the study, the head of the medical institution should be reported the fact and the reason in writing promptly.
- Withdrawal is considered to be in the best interest of the subject or the other subjects.

Subjects may be discontinued from the study for any of the following reasons:

- Lost to follow-up.
- Concomitant treatment with non-authorized medication as defined in the context of this study (in general, any concomitant medication should be discussed with the Contract Research Organization (CRO) Medical Monitor on an ongoing basis).
- If a subject uses any CC or nicotine/tobacco-containing product other than the product/regimen he/she is assigned to, it will be at the discretion of the Investigator to decide whether or not to withdraw the subject from the study.

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

- Non-compliance to the study procedures.

Subjects withdrawn prematurely after randomization will not be replaced and will not be allowed to re-enter the study. All subject withdrawals have to be documented properly in the source documentation and the eCRF.

5.1.4 Violation of Selection Criteria

Subjects who are eligible at Screening, but who do not meet the entry criteria at Admission Day (Day -1) will be considered a screening failure and will be replaced by other subjects.

Subjects who violate the entry criteria prior to enrolment, but who were considered eligible, will be immediately withdrawn from the study when the violation is detected. If subjects are not yet randomized, they can be replaced.

6 INVESTIGATIONAL PRODUCTS AND REFERENCE POINT PRODUCT

6.1 Description

6.1.1 Investigational Products

THS 2.2 Menthol

THS 2.2 Menthol comprises the following components: Menthol Tobacco Stick, Holder, Charger, a Cleaning Tool, a mains power supply, and a USB cable:

Charger: The function of the Charger (Model 4) is to recharge the Holder after use. It contains a battery with sufficient capacity to recharge the Holder approximately 20 times. It is a convenient size to carry around, and can itself be recharged from a mains power source.

THS Tobacco Stick Holder (Holder): The function of the Holder (Model 4.2) is to heat the Menthol Tobacco Stick, delivering an aerosol to the user. The electrical heating is powered from an internal battery which delivers power for about 6 minutes (allowing complete use of a single Menthol Tobacco Stick)

THS Menthol Tobacco Stick (Menthol Tobacco Sticks): The Menthol Tobacco Stick (product code C3 menthol) contains tobacco which, when heated, generates an aerosol. It is custom-designed to be used with the Holder.

The overall objective of the design is to provide an acceptable experience in which the HPHCs level in the aerosol is substantially reduced in comparison with mCCs.

The THS 2.2 Menthol will be provided by the Sponsor.

Per cigarette/Tobacco Stick tar, nicotine, and carbon monoxide yields are normally determined by standardized test methods. The most widely used test method is ISO 4387. PMI has developed a modified version of this method, which improves the determination of tar in products with high water content, which is typical for heated tobacco products ([PMI](#),

2012a; PMI, 2012b; PMI, 2013b). Another method is the more intensive smoking method developed by Health Canada (Health Canada, 1999).

Table 1 below lists the commonly reported measures.

Table 1 Measured aerosol fractions for the THS Menthol Tobacco Sticks

Constituent (mg/THS Menthol Tobacco Stick)	ISO ¹	Health Canada Intense regime ²
Tar/NFDPM	5	12.6
Nicotine	0.5	1.2
Carbon monoxide	1	0.6

¹ International Organization for Standardization ISO machine-smoking regimen. The analytical method has been modified to avoid inaccuracies as a result of condensation from high water-content aerosols.

² Health Canada Intense machine-smoking regimen (55 mL puff volume, 2-second puff duration, 30-second inter-puff interval) (Health Canada, 1999)

Reference Product:

In the study sequences 1 and 2, the reference product to THS 2.2 Menthol is commercially available single brand mCC with a maximum yield of 1 mg nicotine ISO per cigarette.

Menthol CC will not be provided by the Sponsor. All eligible subjects will be asked to purchase their own preferred single-brand mCC prior to Admission. As randomization takes place on Day 0, every study subject needs to buy his/her anticipated amount of single-brand mCC for a total of 2 days plus 2 extra packs.

6.1.2 Reference Point Product

Nicotine replacement therapy gum (Nicorette[®] 2 mg gum) is the reference point product to THS 2.2 Menthol for sequences 3 and 4. The NRT gum will be supplied by the site and reimbursed by the Sponsor.

6.1.3 Packaging and Labeling

At Admission on Day -1, all study subjects will provide the anticipated amount of mCC in sealed packs to the site staff. The cigarette packs provided by the subject should not be opened and the cellophane wrapper should be intact.

Each pack of cigarettes provided by the subject will be labeled to identify which subject the cigarettes belong to (labels should be affixed by the study site trial collaborator to the

cellophane wrapper of the lower part of the pack). Packs of mCC will be labeled to identify necessary information to match the subject with his/hers suppliers.

For the Menthol Tobacco Sticks, the packs and cartons will be pre-labeled with the necessary information including but not limited to product code. The labels will be written in Japanese ([Appendix 4](#)).

6.2 Use of Investigational and Reference Point Products

Subjects will never be requested or forced to smoke and will be free to stop smoking at any time during the study. Subjects caught using any nicotine/tobacco product which is different from the assigned product will be withdrawn from the study. During the Screening period, subjects will be allowed to smoke according to their smoking habits except during the procedures of the Screening Visit [9.1](#) at the discretion of the site.

6.2.1 Admission (Day -1)

Subject will be instructed not to smoke in the morning prior to Admission. Smoking *ad libitum* will be allowed throughout the day of admission except during the procedures until 11:00 PM. All subjects will be allowed to continue smoking *ad libitum* their preferred mCC. All subjects (except women with a positive pregnancy test at Screening or Admission) will undergo a THS 2.2 Menthol test first (using up to 3 Menthol Tobacco Sticks) and subsequently NRT gum test at Day -1 prior to enrolment. Each subject will chew slowly the 2 mg NRT gum for 35 ± 5 minutes while spacing each chew, leading to an administered dose of 1 mg nicotine/product use as per label similarly to the day of single use.

Following agreement that the THS 2.2 Menthol and NRT gum is acceptable, subjects will be enrolled and further randomized to one of 4 treatment sequences using an Interactive Web/Voice Response System.

6.2.2 Investigational Period (Day 0 to Day 3)

During the first washout, each subject will maintain nicotine abstinence from Day -1 at 11:00 PM to the time of single use of his/her allocated product at Day 1. At Day 1, after the single use of the product, subjects will maintain nicotine abstinence for the rest of the day. During the second washout on Day 2, subjects will maintain nicotine abstinence until the time of single use of subject's allocated product at Day 3. Subjects will not be allowed to smoke or use any other nicotine/tobacco-containing products other than the products/regimen they are allocated to.

Time point of first product use during study day (T_0) will be defined as start of the single product use at the single use days. The start time of the use of each product has to be documented on single use days (Day 1 and Day 3). The start of product use for THS 2.2

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

Menthol is defined as the time of the first puff. The start time for mCC corresponds to the lighting of the mCC, and the start time of the NRT is the time of the NRT gum intake. The 30 seconds it takes to pre-heat the Holder will not be taken into account. **The subject must not take a puff of the Menthol Tobacco Stick during the pre-heating time.**

For NRT gum, the start time and the stop time will be documented in appropriate log.

The start of the first product use can be different for each subject both days of product use; however, it must be in the window of 6:00 AM to 9:00 AM.

Single use of products (Day 1 and Day 3):

On Day 1 and Day 3, subjects will use the product they are randomized to only once in the morning between 6:00 AM to 9:00 AM, and will abstain from the product or other nicotine/tobacco-containing items for the rest of the day, i.e., subjects in the THS 2.2 Menthol arm will use one Menthol Tobacco Stick, subjects in the mCC arm will smoke one mCC, and subjects in the NRT gum arm will chew slowly the 2 mg NRT gum for 35 ± 5 minutes while spacing each chew (leading to an administered dose of 1 mg nicotine/product use as per label

	Sequence 1	Sequence 2	Sequence 3	Sequence 4
Day 1	THS 2.2 Menthol	mCC	THS 2.2 Menthol	NRT gum
Day 3	mCC	THS 2.2 Menthol	NRT gum	THS 2.2 Menthol

6.2.3 Day of Discharge/Time of Discharge

On the Day of Discharge (Day 4), smoking will be only allowed after all laboratory procedures and the spirometry have been performed. All examinations of the Day of Discharge will be conducted on Day 4 prior to the time of Discharge.

6.2.4 Safety Period

During the safety follow-up period, subjects are free to smoke according to their usual smoking habits.

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

6.2.5 Stopping Rules for Investigational Product

For safety purposes, smoking the THS 2.2 Menthol or the mCC or the use of the NRT gum should be temporarily stopped in the event of any signs suggesting nicotine overexposure, e.g., gastrointestinal disturbance (nausea, vomiting, diarrhea, stomach, or abdominal pain), cold sweats, headache, dizziness and breathing problems, or any reasons at the discretion of the Investigator.

6.3 Method for Assigning Subjects to Study Arms

Randomization to produce exposure sequence will be done through an Interactive Telephone and Web Response System.

Each sex and each of the smoking level (ISO nicotine levels ≤ 0.6 mg and >0.6 to ≤ 1 mg) will have a quota applied to ensure they represent at least 40% of the total study population allocated within each of the following analysis groups:

- Group 1: composed of sequences 1 and 2.
- Group 2: composed of sequences 3 and 4.

In particular, the maximum number of subjects having the same sex or nicotine level value will be limited to 26 in Group 1 and 10 in Group 2. The randomization of the planned sample size of 62 subjects will be ensured by applying quota to the number of subjects per each sequence (22 subjects for sequences in Group-1, and 9 subjects for sequences in Group-2).

Subjects will be randomly assigned to one of the four product exposure sequences by means of a permuted-block schema. Block size and other randomization details will be available in the randomization plan.

The randomization plan will be generated by an independent statistician and none of the sponsor staff, Investigators or study subjects will have access to the randomization schema prior to randomization.

6.4 Blinding

This is an open-label study; therefore the subjects and investigators will be unblinded to subject's sequence. However, there will be a limited degree of blinding in the data review and data analysis process. In particular, PMI and CRO personnel will be blinded to the randomized sequence as summarized in the following table:

Blinded Study Personnel	End of Blinding Period
PMI and CRO study statisticians	After the Statistical Analysis Plan (SAP) finalization or PMI blind database review ^(*) , whichever comes last.
PMI data manager	After the finalization of PMI blind database review. ^(*)
PMI safety and clinical scientist	After the finalization of PMI blind database review ^(*) . Can be actively un-blinded before that time point in case of the occurrence of any safety question, when appropriate.

(*) As part of the PMI Quality Control (QC) activity, data listings will be reviewed by PMI before database lock, with no access to the randomization sequence information

Any PMI and CRO personnel who are not listed in the above table will be unblinded by default.

6.5 Investigational Product Accountability and Compliance

6.5.1 Dispensing Investigational Product and Reference Point Products

From Day -1 until Day 4, the THS 2.2 Menthol, NRT gum, and mCC will be dispensed by the Investigators or dedicated study collaborator, as per study design. Each dispense of the product will be recorded in a log. The log should include subject number, date and start time of product use. The product will not be promoted for commercial distribution or test market.

6.5.2 Storage and Accountability

The THS 2.2 Menthol, NRT gum and mCC will be stored in a secured site storage place with access limited to the authorized personnel only. Full accountability of the distributed products will be ensured by designated collaborator. Subjects will return each butt of any used Menthol Tobacco Stick or mCC immediately after use to the site study collaborator for accountability. They will also return each used NRT gum after use to the site staff. This will be documented in appropriate log. On the time of discharge, unused mCCs given to the site study collaborator at Admission on Day -1 will be given back to the subjects.

6.5.3 Investigational Product Retention

Unused Menthol Tobacco Sticks and NRT gum will be destroyed if possible, or returned to the Sponsor upon study completion. The Tobacco Heating Devices will be returned to the Sponsor.

6.5.4 Compliance to Investigational Products

Compliance for all arms will be ensured by strict distribution of the products (product by product) and collection of used Menthol Tobacco Sticks, the mCC butts and the NRT gum after use will be documented in appropriate logs.

In addition, for subjects using NRT gum, the compliance will be chemically verified using exhaled CO breath. The cut-off point for the CO breath test value to distinguish tobacco use vs. no tobacco use will be 10 ppm ([Benowitz et al. 2002](#)).

Furthermore, the CO breath test will be considered as one of the measures of compliance during the wash-out days in all subjects.

6.6 Restrictions

6.6.1 Smoking Restrictions

On Day 1 and Day 3, to avoid nicotine cross contamination, users of THS 2.2 Menthol and smokers of mCC will use or smoke, as applicable, in dedicated separate rooms: one room for THS 2.2 Menthol and one room for mCC. Every subject must use or smoke, as applicable alone with an interval between subjects allowing ventilation of the room. Subjects receiving NRT gum must not have access to these rooms.

In the morning prior admission, subjects will be instructed not to smoke. At admission, smoking is only allowed during the designated smoking times from 6:30 AM to 11:00 PM as detailed in the study design. Subjects will not have free access to their NRT gum, mCC or THS 2.2 Menthol, which will be dispensed by the site staff individually as described in [Section 6.5.1](#).

Smoking is not allowed during study procedures except during blood sampling for nicotine PK on Day 1 and Day 3. Furthermore, smoking is not allowed on Day 4 until all laboratory tests and the spirometry have been conducted.

During the days of wash-out or single product use (for mCC and THS 2.2 Menthol arms), no NRT gum or other products supportive to SA must be used or will be provided to the subjects.

6.6.2 Dietary Restrictions

A standard diet will be designed by a dietician for the whole confinement period. For each meal, the caloric and fat content should be controlled in order to avoid a “high-fat” diet. The FDA guidance on food-effect studies for bioequivalency testing identifies a “high-fat” diet as a diet which maintains approximately 50 percent of total caloric content of the meal and is high in calories (approximately 800 to 1000 calories) ([FDA, 2002](#)).

Subjects are not allowed to bring their own food or beverages to the investigational site. Meals will be served according to the schedules provided in [Section 9](#). Additional light snack, fruits, and raw vegetables can be distributed to the subjects without restrictions at any time during confinement as long as they fulfill the above requirements described in this section. Consumption of water is allowed as desired. The same menu and meal schedule will be administered uniformly for all subjects in all study arms.

Fasting state has to be observed for at least 10 hours prior to blood draws for the safety laboratory at the Screening Visit, on Day -1, and Day 4.

6.6.3 Concomitant Medication

For the purpose of this study, no concomitant medication should be taken by the subjects. Any medication with an impact on the CYP2A6 metabolism (as prescription and over-the-counter products) as given below must be avoided as CYP2A6 is involved in the nicotine metabolism (see [Table 2](#)).

In this study the use of hormonal contraception containing estrogens is NOT allowed. This also applies to hormone replacement therapy. Only hormonal contraception with products containing progesterone is allowed during this study. Subjects using estrogens during the study will be withdrawn.

The following drugs and substances are considered as having an impact on CYP2A6 activity ([Lacy et al. 2007](#)). Prior to database close, the concomitant medication will be assessed according to the potential impact on CYP2A6 activity and the potential impact on study results.

Table 2 Drugs and Substances Considered Interacting with CYP2A6

Drug name	Substance Class
Fluoroquinolones, including ciprofloxacin and ofloxacin, nafcillin, rifampicin	Antibiotic
Fluvoxamine, fluoxetine, paroxetine, bupropion,	Antidepressant

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

duloxetine, amitriptyline, imipramine, sertraline, mirtazapine, citalopram, thioridazine	
Haloperidol, perphenazine, chlorpromazine, propoxyphene fluphenazine, clozapine, olanzapine	Neuroleptic
Phenobarbital, primidone, carbamazepine	Antiepileptic
Cholorquine, quinidine	Antirheumatic
Clotrimazole, terbinafine, fluconazole, ketoconazole, miconazole	Antimycotic
Erythromycin, ciprofloxacin, clarithromycin, norfloxacin	Antibiotic
Cimetidine, chlorpheniramine, diphenhydramine, ranitidine	H2-receptor antagonist
Amiodarone, verapamil, mibefradil, mexiletin, propafenone, propranolol, lidocaine	Antiarrhythmic
Losartan, amlodipine, nifedipine	Antihypertensive
Drospirenone, oestrogens	Hormonal contraceptives Agents for hormonal replacement therapy (oestrogens)
Fluvastatin	Cholesterol-lowering agent
Theophylline	Antispasmodic pulmonological agent/Bronchodilator agent
Omeprazole, Lansoprazole	Proton pump inhibitor
Interferon	Antiviral/Immunomodulating agent
Methoxsalen	Anti-psoriatic (substance class Furocoumarins)
Modafinil, Diclofenac, Rofecoxib	Analgesic

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

Insulin	Anti-diabetic
Sildenafil	Phosphodiesterase-Inhibitor (e.g., used for treatment of Erectile dysfunction)
Quinine	Crystalline alkaloid
St. John's Wort	Over-the-counter (herbal remedy) antidepressant
Psoralen	Anti-psoriatic (substance class Furocoumarins)
Pilocapine	Cholinergic agonists (e.g., used for Glaucoma Therapy)

Data sources: [Lacy et al. 2007](#). This list is not exhaustive.

However, the Principal Investigator is responsible for the medical care of the subjects during their participation in this study. Any decisions regarding the prescription of medication will be taken in the best interest of the subject.

If the use of a concomitant medication cannot be avoided for the subject's safety it has to be fully documented (for details, see [Section 7.4.5](#)) Concomitant medications should be followed up with the CRO Medical Monitor on an ongoing basis.

Concomitant medication will first be assessed at Screening Visit. To be eligible for the study any medication with impact on CYP2A6 metabolism must be discontinued at least 14 days prior to Admission to the clinic or for at least five half-lives (whichever is longer). They must not be used during the entire study until the time of discharge. It is at the discretion of the Investigator to assess if a termination of such medication at Screening is medically justified and safe for the subject.

6.6.4 Others

From admission to discharge, practice of intensive exercise and physical work-out will be prohibited as it may impact nicotine absorption profile.

7 STUDY PROCEDURES

Personnel performing study measurements or recording must have the appropriate training fully documented. Quality control (QC) measures have to be in place. All study procedures are provided as an overview in the Schedule of Events ([Appendix 1](#)). In this Section, only the expected/planned time points for the various measurements are given. Considering that not all subjects can have a procedure at the same time point, adequate time windows will be given for each study procedure and each time point in [Section 9](#). Site study collaborator will adhere to the site's Standard Operating Procedures (SOPs) for all activities. Appropriate medical advice will be provided to the subject in case of any medical findings requiring health care.

7.1 Informed Consent

Each subject must give his/her informed consent prior to participating in the study. During the consent process, the Investigator obtaining consent must inform each subject of the nature, risks, benefits of and alternatives to study participation. In addition, each subject must have sufficient time to review the ICF in order to understand it, and have adequate opportunity to ask questions. The ICF must be signed and dated by the subject prior to any study-specific procedures taking place.

7.2 Smoking Cessation Advice and Debriefing

Each subject will be given advice on the risks of smoking 3 times during the study: at the Screening Visit, at Admission (Day -1), and at Day 4. This will take the form of a brief interview according to WHO recommendations ([WHO, 2002](#)). Details of the interview will be recorded in the Source Document File. Information on the risk of smoking will be given to the subjects on an individual basis during a face-to-face meeting between the subject and the Investigator, and may additionally be given in a group session.

In addition to the smoking cessation advice, a debriefing of subjects will be done at each smoking cessation advice session to address any intended or unintended beliefs participants have about the candidate MRTP. The goal of the debriefing would be to help ensure that subjects exit the study with an accurate understanding of product risks, including an understanding that the candidate MRTP has not been demonstrated to be less harmful.

7.3 Support during Smoking Abstinence/Periods of Reduced Smoking

All subjects will be closely monitored by the site staff on Day 0, Day 1; Day 2, and Day 3. This includes monitoring of clinical tests e.g., vital signs, physical examination, and body weight. It also refers to close monitoring of the subject's behavior, AEs, and his/her mood.

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

7.4 Clinical Assessments

Any clinically relevant finding detected during the Screening Visit has to be documented as a concomitant disease. This also applies to clinically relevant findings (e.g., laboratory values, vital signs and ECGs) detected during the Screening Visit. Any untoward medical occurrence in a subject detected during the study which was not present at the Screening Visit must be documented as an AE. Worsening of a pre-existing condition from the Screening Visit onwards will also be documented as an AE. If a clinically relevant finding is detected during the Screening period, the Investigator needs to check if inclusion criterion no. 3 is still fulfilled.

7.4.1 Demographic Data

Demographic data (sex, date of birth/age) will be recorded at the Screening Visit.

7.4.2 Identification of the Current Cigarette Brand

Identification of the current mCC brand(s) smoked by the subject will be done at the Screening Visit and at Day -1. At the Screening Visit, smokers will be asked to bring a pack of their current mCC brand(s) to the site. On Day -1, subjects will hand their mCC supply for the entire confinement period to the site study collaborator. The site study collaborator will document the brand name and yields. A photograph of the front and the side of the cigarette pack supplied by the subject (bearing the tar, and nicotine yields) will be taken by the study site staff in addition to recording the brand name and yields. These photographs will be considered as Source Documentation. A copy of the photographs will be provided to the Sponsor electronically (as Digital Video Disk or Compact Disc [CD]).

7.4.3 Smoking History and Willingness to Quit Smoking

Subjects will be questioned for their smoking history. At Screening and on the Day of Admission (Day -1) this will include questions to evaluate whether the subject has smoked for at least the last 3 consecutive years, to determine the number of CC smoked during the previous 4 weeks, and to evaluate if the CCs smoked during the previous 4 weeks were mCCs. At the Screening Visit only, the subject will also be asked if he/she plans to quit smoking within the next 3 months. In addition, the subject will be asked if he/she has used nicotine-containing products other than commercially available CC (either tobacco-based products or nicotine replacement therapy), electronic cigarettes or similar devices, within 4 weeks prior to assessment.

At Screening and on the Day of Admission (Day -1), subjects will also be asked if they are willing to abstain from smoking for at least 4 days (as required in the study protocol inclusion criteria). Only subjects who are prepared and able to comply with this requirement will be considered for participation in the study.

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

7.4.4 Demonstration and Trial of the THS 2.2 Menthol and NRT Gum

All subjects will have a demonstration of the THS 2.2 Menthol product and the NRT gum at the Screening Visit by the study site collaborator. On Day -1 as last procedure of the eligibility assessments on that day, subjects will have a product test prior to enrolment, first THS 2.2 Menthol (use of up to three Menthol Tobacco Sticks) and subsequently NRT gum. Each subject will chew slowly the 2 mg NRT gum for 35±5 minutes while spacing each chew, leading to an administered dose of 1 mg nicotine/product use as per label similarly to the day of single use.

In female subjects, the THS 2.2 Menthol and NRT gum product tests must only be done after pregnancy is excluded by a urine pregnancy test. Only subjects willing and ready to use the THS 2.2 Menthol and NRT gum can be enrolled into the study.

7.4.5 Medical History, Concomitant Disease, Previous and Ongoing Medications

Relevant medical history and any concomitant disease will be documented at the Screening Visit. Medical history is defined as any condition that started prior to and ended prior to Screening. A concomitant disease is defined as any condition that started prior to the Screening Visit and is still ongoing at the Screening Visit.

Prior medication taken within 4 weeks prior to Screening Visit and any concomitant medication needs to be documented. Any medication which was started prior to the Screening Visit and is still being taken by the subject will be considered a concomitant medication. Medication initiated after Screening is also referred to as concomitant medication. This applies to both prescription and over-the-counter products.

Records of any medication taken must include the drug name (preferably both generic and trade name), route of administration (e.g., oral, intravenous), total daily dose/unit (e.g., expressed in mg, ml, or IU), indication, the start and (if applicable) the stop date (day, month, and year). Any therapy changes (including changes of regimen) during the study are to be documented. Any concomitant medication that is still being taken by the subject at EOS will be recorded on the eCRF.

7.4.6 Physical Examination

A physical examination will be conducted at the Screening Visit, at Admission (Day -1), and at the Day of Discharge (Day 4).

7.4.7 Body Height and Weight

Body weight will be recorded at the Screening Visit, at Admission (Day -1) and at the Day of Discharge (Day 4). Body height will be measured at the Screening Visit, only. The BMI will

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

be calculated from the body weight and height using the following formula, rounded to one decimal place:

$$\text{BMI} = \frac{\text{weight in kilograms}}{\text{height in meters}^2} = \frac{\text{kg}}{\text{m}^2}$$

7.4.8 Vital Signs

Systolic and diastolic blood pressure, pulse rate, and respiratory rate will be measured at the Screening Visit, at Admission, and at every day in confinement. All parameters will be recorded in supine position after the subject has rested for at least 5 minutes.

For every measurement, it will be documented if the subject has smoked within 15 minutes prior to the measurement.

7.4.9 Other Clinical Assessments

7.4.9.1 Spirometry

Spirometry with and without a short-acting bronchodilator will be done at the Screening Visit to evaluate inclusion/exclusion criteria (the post-bronchodilator results). At screening, spirometry without bronchodilator will be done first, and then, spirometry with bronchodilator. Furthermore, spirometry without bronchodilator will be performed at Day -1 (baseline values) and at Day 4 (for comparison with the baseline values). Spirometry has to be done prior to smoking the first cigarette of the day.

Spirometry will follow the 2005 testing and quality recommendations by the American Thoracic Society/European Respiratory Society Joint Task Force on the standardization of spirometry along with the electronic data submission and documentation processes. Spirometry predicted values will be standardized to the NHANES III predicted set.

All personnel performing lung function testing should have the appropriate training and quality control (QC) measures should be put into place and be properly documented and filed at the pulmonary function laboratory (including the records of the calibration, if applicable). The FEV₁ and FVC will be recorded.

The subject will be submitted to a spirometry with maximum voluntary ventilation measurement.

For spirometry, assessed parameters will include:

- FEV₁.
- FVC.

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

- FEV₁/FVC.

7.4.9.2 Electrocardiogram

An ECG will be recorded at Screening and on the following study days: Day 1 and Day 3. The ECG testing will be performed as per the site local practice. A standard 12-lead ECG will be recorded after the subject has rested for at least 10 minutes in supine position.

The following parameters will be documented: heart rate, PR interval, QRS interval, QT interval, and QTc interval corrected by the ECG device according to Bazett's formula. Every ECG has to be assessed as normal, abnormal – clinically not relevant, or abnormal – clinically relevant. A diagnosis has to be provided in the Case Report Form for all ECGs assessed as abnormal – clinically relevant. Electrocardiogram print-outs will be interpreted by a qualified physician. Any print-outs of ECGs on thermo-sensitive paper must be photocopied, initialed, dated and stapled together for inclusion in the Source Data File.

7.4.9.3 Chest X-ray

A chest X-ray (anterior-posterior and left lateral views) will be assessed during the Screening period to exclude subjects with relevant pulmonary diseases. Subjects will be referred to a radiology facility for this procedure. No new examination is required if the subject can present a chest X-ray with anterior-posterior and left lateral views at the Screening Visit which is not older than 6 months.

7.5 Biomarker Assessment

All bioanalytical assays and laboratory assessments ([Section 7.6](#)) will be carried out using validated methods. The bioanalytical methods used will be documented in the Bioanalytical Plans/Report. A list of laboratories is provided in [Appendix 2](#).

The start time of the use of each product has to be documented on single use days (Day 1 and Day 3). For NRT gum specifically, the time of NRT gum use (start time T₀), and the stop time of NRT gum use needs to be recorded.

Precautions should be taken during blood sampling and processing to prevent the contamination of samples with environmental nicotine or CO.

7.5.1 Biomarker of Exposure

7.5.1.1 Biomarkers of Exposure to CO and COHb

Carboxyhemoglobin measured in blood and exhaled CO will be investigated as a measure of exposure to CO. The CO breath test will also serve as a measure of compliance in subjects using NRT gum as well as on the wash-out days in all subjects.

CO Breath Test

CO in exhaled breath will be measured using the Smokerlyzer[®] device such as Micro+[™] Smokerlyzer[®] device or similar. The test will be performed for all subjects including the subjects using the NRT gum.

A CO breath test will be conducted once on Day -1 and Day 4:

On Day 0, Day 1, Day 2, and Day 3, four CO breath tests will be done per day. On Day 1 and Day 3, the first test per day will be performed within 15 minutes prior to T₀. The three other test will be conducted as described in [section 9](#). On the wash-out days (Day 0 and Day 2), the first test will be conducted between 8:00 AM-9:30 AM. The three other tests will be conducted as described in [section 9](#).

Carboxyhemoglobin

Tests for COHb measurement will be performed at a local laboratory.

Blood samples will be taken as follows at Day 1 and Day 3:

A total of 5 blood samples will be taken. The first sample will be taken within 15 minutes prior to using the first product (T₀). Thereafter, the sampling times in relation to T₀ are at 15 minutes, 60 minutes, 4 hours and 12 hours post-T₀.

7.5.1.2 Biomarkers of Exposure to Nicotine

Blood samples to measure nicotine in plasma will be taken as follows:

Single Use on Day 1 and Day 3 for THS 2.2 Menthol and mCC only:

A total of 16 blood samples will be taken for a 24-hour profile (Day 1 and Day 3). The first blood sample will be taken within 15 minutes prior to the single use (T₀). Times of sampling are thereafter in relation to T₀: T₁ after 2 minutes, T₂ after 4 minutes, T₃ after 6 minutes, T₄ after 8 minutes, T₅ after 10 minutes, T₆ after 15 minutes, T₇ after 30 minutes, T₈ after 45 minutes, T₉ after 60 minutes, T₁₀ after 2 hours, T₁₁ after 4 hours, T₁₂ after 6 hours, T₁₃

after 9 hours, T₁₄ after 12 hours and T₁₅ after 24 hours (this sample will be drawn during the day following product use, i.e., wash-out).

Single Use on Day 1 and Day 3 for NRT gum only:

A total of 16 blood samples will be taken for a 24-hour profile (Day 1 and Day 3). The first blood sample will be taken within 15 minutes prior to the single use (T₀). Times of sampling are thereafter in relation to T₀: T₁' after 10 minutes, T₂' after 20 minutes, T₃' after 25 minutes, T₄' after 30 minutes, T₅' after 35 minutes; T₆' after 40 minutes; T₇' after 45 minutes, T₈' after 60 minutes, T₉' after 2 hours, T₁₀' after 3 hours, T₁₁' after 4 hours, T₁₂' after 6 hours, T₁₃' after 9 hours T₁₄' after 12 hours and T₁₅' after 24 hours (this sample will be drawn during the day following product use, i.e., wash-out).

7.5.2 CYP2A6 Activity

CYP2A6 activity will be measured in plasma on Day -1. CYP2A6 activity drives the metabolism of nicotine to cotinine and subsequent metabolites. In this study the CYP2A6 activity will be measured using the metabolic molar ratio of *trans*-3'-hydroxycotinine/cotinine (Jacob et al. 2011).

7.6 Laboratory Assessments

A list of laboratories is provided in [Appendix 2](#).

7.6.1 Clinical Chemistry, Hematology, and Urine Analysis for Safety Panel

Hematology, clinical chemistry and urine analysis for the safety panel will be measured at Screening, at Day of Admission (Day -1), and at the Day of Discharge (Day 4). Tests will be conducted at a local laboratory or the site. Blood will be taken after no less than the 10 hours of fasting (see [Section 6.6.2](#)). The urine test will be performed semi-quantitatively as urine test at the site. Parameters to be measured are listed in [Table 3](#).

Table 3. Clinical Laboratory Parameters for Safety Panel

Hematology	Clinical chemistry	Urine analysis
- Hematocrit	- Albumin	- pH
- Hemoglobin	- Total protein	- Bilirubin
- Mean corpuscular hemoglobin	- Alkaline phosphatase	- Glucose
- Mean corpuscular hemoglobin concentration	- Alanine aminotransferase	- Nitrite
- Mean corpuscular volume	- Aspartate aminotransferase	- Red blood cell traces
- Platelet count	- Blood urea nitrogen	- Protein
- Red blood cell count	- Creatinine	- Specific gravity
- White blood cell (count) (WBC)	- Gamma-glutamyl transferase	
- Differential WBC count:	- Fasting glucose	
• Neutrophils	- Lactate dehydrogenase	
• Basophils	- Potassium	
• Eosinophils	- Sodium	
• Lymphocytes	- Total bilirubin	
• Monocytes	- Direct bilirubin	
	- Total cholesterol	
	- Triglycerides	

7.6.2 Serology

A test for Hepatitis B surface antigen, Hepatitis C virus and HIV (anti-HIV1/2 and p24 antigen) will be done at Screening. In case of positive results, the subject will be referred to appropriate medical care.

7.6.3 Urine Drug Screen

A urine drug screen will be performed at the site at the Screening Visit and at the day of Admission. The urine will be screened for amphetamines, barbiturates, benzodiazepines, cannabinoids, cocaine, and opiates.

7.6.4 Urine Cotinine Screening

A urine cotinine test will be performed at Screening and at Admission to the clinic in order to confirm the subject's smoking status. The test must detect cotinine with a cotinine of ≥ 200 ng/mL.

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

7.6.5 Alcohol Breath Test

Subjects will undertake a breath alcohol test at the Screening Visit and at Admission to the clinic using a newly calibrated alcometer device.

7.6.6 Urine Pregnancy Testing

All female subjects will undergo pregnancy testing at the Screening Visit, at Admission to the clinic, and at the Day of Discharge (Day 4). Female subjects with a positive pregnancy test at the Screening Visit or on Day -1 cannot be enrolled and are considered a screening failure. Pregnancy in such subjects will not be followed up as no exposure to the THS 2.2 Menthol will have occurred. Product test at Admission must be done only in female subjects with a negative pregnancy test. In any case of a positive pregnancy test, the Investigator will inform the subject about the risks associated with smoking during pregnancy. In the event of unclear urine pregnancy test in peri-menopausal women, absence of pregnancy should be confirmed by a serum follicle stimulating hormone level >20 IU/l.

All pregnancies detected during the study must be reported and handled as described in [Section 8.5.1](#).

7.7 Sample Handling, Storage, and Shipment

All blood samples are to be tested at a central laboratory with the exception of COHb blood samples and the safety laboratory samples which will be tested at a local laboratory (see [Appendix 2](#)). The urine drug screen, urine pregnancy tests and urine cotinine tests will be done by the site study collaborator at the site. The tests will be provided by the sites.

Detailed procedures for handling of samples are described in the separate sample handling manual (SHM). Safety laboratory samples will be destroyed as by the laboratories standard procedures. All other samples will be destroyed once the Clinical Study Report (CSR) has been finalized. The facility/-ies at which the samples are stored will be informed in writing by the Sponsor when destruction of the samples shall be performed.

7.7.1 Blood samples

Blood samples will be drawn by qualified and trained site personnel. Subjects should be in a seated position during blood collection. The maximal total volume of blood drawn for each subject will be around 190 mL. The required aliquots and volumes for assessments of blood/plasma parameters and tests are summarized in the SHM.

7.7.2 Urine samples

Spot urine samples will be taken for urine drug screen, cotinine screen, pregnancy tests and safety urinalysis.

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

7.8 Questionnaires

The subject questionnaires and the Visual Analogue Scale (VAS) will be entered by the subject directly in the electronic patient reported outcomes (ePRO) device. The questionnaires and the VAS will be reviewed for completeness by the study site collaborator and subjects will be requested to complete any missing information.

Symptoms or worsening of symptoms as documented on any of the questionnaires or the VAS do not need to be documented as AEs because the questionnaire and the VAS will be analyzed as part of the report. However, it is at the discretion of the Investigator to document such symptoms also as AEs. The main source for AE collection will be the face-to-face interview between the subject and site study collaborator using, open, non-directive questions (see [Section 8](#)).

7.8.1 Fagerström Test for Nicotine Dependence (revised version)

Potential nicotine dependence will be assessed at Screening using the FTND in its revised version ([Heatherton et al, 1991](#)), as updated in 2012 ([Fagerström et al., 2012](#)).

The questionnaire consists of 6 questions which have to be answered by the subject himself/herself. The scores obtained on the test permit the classification of nicotine dependence into three levels: Mild (0 to 3 points); Moderate (4 to 6 points); Severe (7 to 10 points) ([Fagerström et al., 2012](#)).

7.8.2 Assessment of Cough

Subjects will be asked if they have experienced a regular need to cough e.g., coughing several times in the last 24 hours prior to assessment. If the answer is 'yes', they will be asked to complete a VAS, 3 Likert scales and an open question also assessing the previous 24 hours. The questionnaire will be asked prior to product use.

The VAS will assess how bothersome cough is to the subject ranging from 'not bothering me at all' to 'extremely bothersome'.

Furthermore, subjects will be asked to assess the intensity and frequency of cough and the amount of sputum production on Likert scales:

- The intensity of cough will be assessed on a 5-point Likert scale ranging from 1 to 5:
1 = very mild; 2 = mild; 3 = moderate; 4 = severe; 5 = very severe.
- The frequency of cough will be assessed on a 5-point Likert scale ranging from 1 to 5:
1 = rarely; 2 = sometimes; 3 = fairly often; 4 = often; 5 = almost always.

- The amount of sputum production will be assessed on a 4-point Likert scale ranging from 0 to 3: 0 = no sputum; 1 = a moderate amount of sputum; 2 = a larger amount of sputum; 3 = a very large amount of sputum.

Finally, subjects will be asked with an open question if there are any other important observations that they would like to share with the staff about their coughing.

Assessments will be done on a daily basis from Day 0 to Day 4. On Day 2 and Day 4, questionnaire must be asked 24 hours after T₀ of Day 1 and after 24 hours after T₀ of Day 3.

7.8.3 Modified Cigarette Evaluation Questionnaire (modified version)

Product evaluation will be assessed using the MCEQ ([Cappelleri et al., 2007](#)). The MCEQ assesses the degree to which subjects experience the reinforcing effects of smoking, by measuring:

- Smoking satisfaction (satisfying, tastes good, enjoy smoking).
- Psychological rewards (calms down, more awake, less irritable, helps concentrate, reduces hunger).
- Aversion (dizziness, nauseous).
- Enjoyment of respiratory tract sensations (single-item assessment).
- Craving reduction (single-item assessment).

This questionnaire will only be completed by the subjects who use the THS 2.2 Menthol or smoke mCC during the study in sequence 1 and 2. The MCEQ will be completed by subjects on Day 1 and Day 3.

7.8.4 Questionnaire of Smoking Urges brief

To assess the urge-to-smoke, all subjects will be asked to complete a 10-item brief version of the QSU-brief ([Cox et al. 2001](#)). The QSU-brief is a self-reported questionnaire with 10 items to be rated on a 7-point scale, ranging from 1 (strongly disagree) to 7 (strongly agree). Higher scores in this questionnaire indicate a higher urge to smoke.

The findings in this brief version were consistent with the expressions of craving found in the 32-item version of the QSU-brief ([Tiffany et al. 1991](#)). The findings supported a multi-dimensional conceptualization of craving to smoke and demonstrated the utility of a brief multi-dimensional measure of craving ([Cox et al. 2001](#)).

The QSU-brief will be completed by the subject himself/herself at single use study days.

7.8.4.1 THS 2.2 Menthol and CC arms QSU-timing:

For subject on THS 2.2 Menthol or on mCC: first sample within 15 min prior to T_0 , 9 assessments thereafter in relation to T_0 : 15 min, 30 min, 45 min, 60 min, 2 hr, 4 hrs, 6 hrs, 9 hrs, and 12 hrs after T_0 .

7.8.4.2 NRT gum arm QSU-timing:

For subjects on NRT gum: first sample within 15 min prior to T_0 , 9 assessments thereafter in relation to T_0 : 20 min, 30 min, 45 min, 60 min, 2 hr, 4 hr, 6 hrs, 9 hrs, and 12 hrs after T_0 .

8 ADVERSE EVENTS

8.1 Definitions

8.1.1 Adverse Events

The FDA MRTP guidelines specify the following definition for AEs for tobacco products ([FDA, 2012a](#)):

An AE is any health-related event associated with the use of a tobacco product in humans, which is adverse or unfavorable, whether or not it is considered related to the tobacco product, as defined by the MRTP guidelines.

An AE is defined as any untoward medical occurrence in a subject, or clinical investigation subject administered an IP, which does not necessarily have a causal relationship with the IP or reference point product. An AE can therefore be any unfavorable and unintended sign (including a clinically relevant abnormal laboratory finding), symptom, or disease temporally associated with the use of an IP or reference point product, whether or not it is considered related to the IP or reference point product.

8.1.2 Serious Adverse Events

An SAE is defined as, but not limited to, any untoward medical occurrence that:

- Results in death.
- Is life-threatening.
- Requires inpatient hospitalization or prolongation of existing hospitalization.
- Results in persistent or significant disability/incapacity, or
- Is a congenital anomaly/birth defect.

Important medical events that may not result in death, be life-threatening, or require hospitalization may be considered an SAE when, based on appropriate medical judgment, they may jeopardize the subject or the subject may require medical or surgical intervention to prevent one of the outcomes listed in the above definitions.

Any pre-planned hospitalizations that are known at the time of signing the ICF will not be recorded as SAEs; however, they will be recorded as AEs only. Any AE that occurs during this pre-planned hospitalization will be considered according to the above definitions.

8.2 Assessment of Adverse Events

The Investigator is responsible for obtaining, assessing and documenting all AEs during the study.

8.2.1 Collection of Information

Adverse event information will be collected from the time of signature of the ICF onwards until EOS either by the Investigator via spontaneous reporting or by the use of consistent, open, non-directive questions from study site collaborator (e.g., “Have you had any health problems since the previous visit/How are you feeling since you were last asked?”). At the discretion of the Investigator, the collection of AE information may also be triggered from his/her review of the subject questionnaires and the VAS. However, the main source for AE collection will be the face-to-face interview with the subject.

Information recorded will include: verbatim description of the AE, start and stop dates and times, seriousness, severity (intensity), action taken (e.g. whether or not the AE led to the subject’s withdrawal from the study), and outcome (e.g. resolved, withdrawal due to AE).

For each AE the intensity will be graded on a 3-point intensity scale (mild, moderate, severe) using the definitions provided in [Section 8.2.3](#).

Any exacerbation/worsening or increased frequency of an AE or pre-existing condition shall be evaluated and recorded.

Correct medical terminology/concepts are preferred when recording AE terms, and abbreviations must be avoided. Wherever possible, a diagnosis is to be used to describe an AE rather than individual signs and symptoms (e.g. record ‘pneumonia’ rather than ‘fever’, ‘cough’, ‘pulmonary infiltrate’ or ‘septicemia’ rather than ‘fever’ and ‘hypotension’ following blood sample).

Any AE that meets the serious criteria must be recorded both on the AE report form of the eCRF and on a separate SAE report form (see [Section 8.2.3](#)).

8.2.2 Period of Collection

8.2.2.1 Screening Period

All existing health conditions identified during the Screening period will be recorded as concomitant disease and the subject’s eligibility for admission to the study will be reviewed. Any AEs which occur during the screening period will be captured by the study site collaborator and assessed by the Investigator in order to establish relationship or relatedness in respect to study procedures. Only the study procedures-related AEs will be reported in the CSR and in accordance with respective regulatory guidelines.

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

8.2.2.2 Admission Day until the End of Study

From Admission onwards until Day of Discharge, all AEs will be actively collected by the study site collaborator.

Any new, clinically relevant, abnormal finding or worsening of a pre-existing condition/concomitant disease detected during the study will be documented as an AE and/or SAE as described in the safety management plan.

During the safety follow-up period new AEs and/or SAEs will be recorded after spontaneous reporting by the subject. Serious AEs will be reported by the Investigator as described in this document and the Safety Management Plan. Any ongoing AEs/SAEs during the safety follow-up period will be actively followed up by the site until they have been resolved, stabilized (i.e., no worsening of condition), or an acceptable explanation has been found.

At the end of the safety follow-up period all ongoing AEs/SAEs will be followed up by the Investigator on behalf of the Sponsor (see [Section 8.3](#)) until they have resolved, stabilized (i.e., no worsening of condition), or an acceptable explanation has been found.

8.2.3 Intensity of Adverse Event

For each AE, the intensity will be graded by the Investigator on a 3-point intensity scale (mild, moderate, severe) using the following definitions:

- **Mild:** The AE is easily tolerated and does not interfere with activities of daily living (ADL).
- **Moderate:** The AE interferes with ADL, but the subject is still able to function.
- **Severe:** The AE is incapacitating and requires medical intervention.

8.2.4 Relationship to Investigational Product and Relationship to Study Procedures

According to the Council for International Organizations of Medical Sciences VI Working Group, there are no definitive methods for distinguishing most adverse drug reactions (i.e., events that are causally attributed to the IP or reference product) from clinical AEs that occur as background findings in the population and have only temporal association with the IP or reference product.

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

In general, all AEs and/or SAEs will be assessed by the Investigator as either ‘related’ or ‘not related’ to IP or reference product as described below. In addition to the assessment of the relationship of the clinical event to the IP or reference product, the Investigator shall document a potential relationship of the clinical event to any particular study procedure.

- **Not related:** The temporal relationship of the clinical event to IP administration or a study procedure makes a causal relationship unlikely, or, concomitant medication, therapeutic interventions, or underlying conditions provide a sufficient explanation for the observed event.
- **Related:** The temporal relationship of the clinical event to study IP administration or a certain study procedure makes a causal relationship possible, and concomitant medication, therapeutic interventions, or underlying conditions do not provide a sufficient explanation for the observed event.

8.2.5 Expectedness

An AE will be regarded as ‘unexpected’ if its nature or severity is not consistent with information already known about the IP or reference product, and/or has not been previously observed and is not listed in the current IB. The IB provides further detail on signs or symptoms that might be expected with the use of the IP, including information relating to device malfunction or misuse.

NRT gum-related AEs are listed on the product label.

8.3 Reporting of Serious Adverse Events

Any SAEs reported or observed during the study after signature of the ICF until the end of the safety follow-up period (i.e. up to 7 days after study Discharge), whether or not attributable to the IP, to any other medication or to any study procedures, or any SAE related to the product and spontaneously reported after the safety follow-up must be reported by the Principal Investigator **within 24 hours after first awareness by any party** to United BioSource Corporation (UBC) Safety and to the Sponsor, and the head of the Investigational site.

An SAE report form must be faxed or e-mailed as attachment to:

UBC Safety:	Phone number: +41 22 596 4444
	E-mail: EUsafety@unitedbiosource.com
	Address: United BioSource Corporation 16, Chemin des Coquelicots

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

1214 Vernier/Geneva
Switzerland

Sponsor Contact:	Mobile:	+41 (79) 484 9068
Tamara Koval	E-mail:	Tamara.Koval@pmi.com
	Address:	Philip Morris Products S.A. R&D Innovation Cube 5 Quai Jeanrenaud 2000 Neuchâtel Switzerland

The Principal Investigator/the head of the investigational site is responsible for local reporting (e.g., to the IRB) of SAEs that occur during the study, according to local regulations.

Any follow-up information will be detailed in a subsequent SAE report form and sent to UBC Safety and the Sponsor

The follow-up SAE report form must include the minimum information required for form completion and only changed/new information needs to be specified. Information provided in the follow-up SAE report form supersedes any information that is initially reported.

All SAEs will be followed-up by the Investigator until resolution or until the Investigator considers the event to be stabilized (i.e., no worsening of condition), or an acceptable explanation has been found (e.g., chronic condition).

The SAE report form to be used in this study is provided as a separate document. All SAEs will be recorded on the relevant eCRF page, in addition to the SAE report form.

8.4 Reporting of Other Events Critical to Safety Evaluations

8.4.1 Abnormal Results of Laboratory Tests

Any clinical safety laboratory test result that is outside of the normal reference range will be reviewed by the Investigator and assessed for clinical relevance. If the Investigator considers the abnormal result to be of clinical relevance, then it must be recorded as a concomitant disease at Screening, or if not present at Screening, as an AE during the study.

The grading scheme shown in (reference to the Common Terminology Criteria for Adverse Events and Common Toxicity Criteria [CTCAE] version 4.03) will be used by the Investigator to assess abnormal laboratory AEs as follows:

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

- All Grade 1 abnormal laboratory values will be evaluated by the Investigator with respect to baseline value and clinical relevance. If considered to be clinically relevant, the Investigator must report it as an AE. All Grade 2 and higher abnormal laboratory values must be reported as or linked to an AE/concomitant disease.
- If a subject has Grade 2 and higher abnormal laboratory values at Screening, it is at the discretion of the Investigator to enroll the subject or not. This decision must be documented in the source documentation and captured in the eCRF.
- If there is any worsening in grade from Grade 2 and above during the study, the Investigator must report this worsening as an AE.
- Where there is no grading available, the abnormal laboratory value will be evaluated by the Investigator, and assessed for clinical relevance. If considered to be clinically relevant, the Investigator will report it as an AE.
- Any other abnormal clinical laboratory result (including those that are not part of the core safety assessments) can, at the discretion of the Investigator, be reviewed and assessed. Even if they do not meet the criteria of the CTCAE grading scheme (please see above), the Investigator may consider them to be of clinical relevance and, if they are, must report them as AEs.
- In general, laboratory values will be recorded as ‘increased <lab parameter>’ or ‘decreased <lab parameter>’ to ensure consistency of recording/coding.

All other information (e.g., relationship to IP, intensity, seriousness, outcome) will be assessed as for other AEs.

8.4.2 Abnormal Results of Others Tests or Investigations

An ongoing medical condition or clinically relevant finding detected during the Screening Visit (including elevated laboratory parameters), will be considered a concomitant disease and the subject’s eligibility for admission to the study will be reviewed.

Any new, clinically relevant, abnormal finding or worsening of a pre-existing condition detected during the study after Screening until the EOS will be documented as an AE.

Any new onset of symptoms or worsening of pre-existing symptoms identified through the study questionnaires may be documented as AEs at the discretion of the Investigator.

8.5 Actions and Follow-up

8.5.1 Reporting and Follow-up of Pregnancies

For pregnancies detected during the Screening period and prior to first THS 2.2 Menthol use, the subject will be considered as a screening failure and removed from the study. No Pregnancy Form will be filled; however, the diagnosed pregnancy must be captured in the Screen Failure eCRF.

All pregnancies occurring after signature of the ICF and diagnosed after first exposure to the IP until completion of the study must be reported by the Investigator.

Any pregnancy potentially associated to exposure to the IP, including pregnancies spontaneously reported to the Investigator after the EOS must be reported by the Investigator and followed-up. Potential association with exposure to the IP is defined as the conception date being calculated before the last exposure to the IP.

The Investigator will complete a Pregnancy Form (provided as a separate document) for all pregnancies diagnosed (including positive urine pregnancy tests).

The procedure to report a pregnancy and provide any additional/follow-up information to UBC Safety and the Sponsor must be followed in the same manner and within the same timelines as described for an SAE (see [Section 8.3](#)). In addition, each pregnancy has to be reported as a non-serious AE. No invasive procedures, including drawing of blood, must be done in such subjects after the discovery of pregnancy.

United BioSource Corporation Safety will follow-up pregnancies only if they were detected after first product use (i.e., after THS 2.2 Menthol test on Admission Day). If pregnancies are to be followed-up, they will be followed-up until an outcome is reached (e.g., normal delivery, spontaneous abortion, or voluntary termination). Any pregnancy complications, adverse pregnancy outcomes, or maternal complications will be recorded.

The Principal Investigator/head of the investigational site is responsible for informing the IRB of any pregnancy that occurs during the study and its outcome, according to local regulations.

8.5.2 Adverse Events Leading to Withdrawal

Subjects who are withdrawn from the study because of an AE will undergo the procedures, as described for the Day of Discharge, as soon as possible and will enter the period of safety follow-up. The Investigator will follow-up these AEs until they have resolved, stabilized (i.e., no worsening of condition), or an acceptable explanation has been found.

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

8.5.3 Investigational Device Misuse

Any occurrences of the THS Tobacco Stick Holder or THS Charger misuse (use not in accordance with its label and instruction) by a subject, will be documented by the Investigator or his/her designated study collaborator using a Device Issue Log.

Investigational device misuse may result in use-related hazards.

Use-related hazards are derived from the US Food and Drug Administration Medical Device Use-Safety Guidance ([FDA, 2012b](#)):

- Hazards caused specifically by how a device is used
- Unanticipated use scenarios (e.g., modification of Charger, applying any chemicals, using conventional cigarettes, mechanical damage of the device, etc.) that result in hazards must be documented and reported by the Investigator.

According to FDA Device Regulation, data should be collected regarding the use-related hazards that have occurred with device and when information pertaining to device use safety is extensive, it is helpful to provide it in summary form that highlights the most important issues, considerations, resolutions, and conclusions. The level of detail of device use documentation submitted should be consistent with the level of concern of use-related hazards for the device.

Furthermore, any misuse of the THS Tobacco Stick Holder or THS Charger, that lead to an AE/SAE will follow the same processes as described above.

8.5.4 Investigational Device Malfunction

Any occurrences of malfunction of the Tobacco Stick Holder or Charger will be documented by the Investigator or his/her designated study collaborator using a Device Issue Log.

Furthermore, any malfunctions of the Tobacco Stick Holder or Charger that lead to an AE/SAE will follow the same processes as described above.

9 STUDY ACTIVITIES

A detailed schedule of assessments can be found in [Appendix 1](#). The time points shown are to be considered the time of assessment for the first subject. As not all subjects can be treated at the same time, a short time window will be implemented for subsequent subjects. Measurements not conducted at the exact time point, but conducted within the given time window (if applicable) do not constitute a protocol deviation but an accepted variability for the given time point.

In general, if no start time for the procedure is provided, then the procedure can be performed at any time during the day.

9.1 Screening Visit

The Screening Visit will be performed within 4 weeks (Day -29 to Day -2) prior to Admission (Day -1). Subjects will attend the investigational site in at least a 10-hour fasting state for clinical laboratory to be assessed.

The following assessments will be performed at the Screening Visit ([Table 4](#)) (the sequence of the assessment will be at the discretion of the site but all of them must be done after signature of the ICF).

Table 4 Time Schedule – Screening Visit

Time	Blood sample	Procedures	Additional information
Start of procedure		Informed consent/Subject Information Sheet	
		Demographic data	On the day of ICF signature
		Advice on the risks of smoking and debriefing	
		Smoking history	
		Willingness to abstain from smoking in the next 3 months	
		Readiness to accept interruptions of smoking for up to 4 days	
		FTND questionnaire	

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

	Prior medication (4 weeks prior to Screening visit)and concomitant medication	
	Medical history/concomitant diseases	
	Vital signs (pulse rate, systolic and diastolic blood pressure, respiratory rate)	At least 5 min in supine position prior to measurement
	Height, weight, including calculated BMI	
√	Clinical laboratory parameters (hematology, urine analysis, clinical chemistry)	At least in 10 hours fasting condition
√	Serology for HIV and Hepatitis B and C	
	Identification of current mCC brand	
	Urine drug screen	
	Alcohol breath test	
	Urine pregnancy test for all female subjects	
	THS 2.2 Menthol and NRT gum product demonstration	
	Spirometry without short-acting bronchodilator first, and then with	Has to be done at least 1 hour after smoking
	ECG	At least 10 min in supine position prior to recording
	AE/SAE questioning	If the Screening Visit is performed on two separate days the AE/SAE questions will be asked again
	Physical examination	
	Chest X-ray (if not performed 6 months prior to Screening)	
	Urine cotinine screening test	
	Inclusion/exclusion criteria	

Abbreviations: AE = Adverse event; BMI = Body mass index; mCC = menthol conventional cigarette(s); ECG = Electrocardiogram; FTND = Fagerström Test for Nicotine Dependence (revised version); HIV = Human immunodeficiency virus; NRT gum = nicotine replacement therapy gum; SAE = Serious adverse event; THS = Tobacco Heating System

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

9.2 Admission

The following assessments will be performed at Admission (Day -1) ([Table 5](#)):

Table 5. Time Schedule – Day -1 Admission

Time	Blood sample	Procedures	Additional information
Start of procedure			
		AE/SAE recording, concomitant medication	All day
		Advice on the risks of smoking and debriefing	
		Readiness to accept interruptions of smoking for up to 4 days	
	√	<i>trans</i> -3'-hydroxycotinine and cotinine (CYP2A6 activity) in plasma	The subject should not have smoked in the morning until this assessment Must be done prior to smoking
		Spirometry	Must be done prior smoking
6:30 AM		Beginning of smoking period	
	√	Clinical laboratory parameters (hematology, urine analysis, clinical chemistry)	To be done at least in 10-hour fasting condition
Prior to 10:00 AM		Breakfast	
		Urine pregnancy test for all female subjects	
Prior to 11:30 AM		Urine cotinine screening test	
Prior to 11:30 AM		Urine drug screen	
Prior to 11:30 AM		Vital signs	At least 5 min in supine position prior to measurement
Prior to 11:30 AM		Physical examination, weight and calculated BMI	
Prior to 11:30 AM		Identification of current mCC brand	

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

Philip Morris Products S.A.
ZRHM-PK-05-JP

Clinical Study Protocol
Final 1.0 / 21 June 2013

Confidential
Page 74 of 124

Prior to 11:30 AM	Smoking history	
Prior to 11:30 AM	Alcohol breath test	
Prior to 11:30 AM	CO breath test	
Prior to 2:30 PM	Lunch	
Afternoon	Snacks	
	Product test for THS 2.2 Menthol and NRT	The THS 2.2 Menthol test should be done first and then the NRT gum test. Subject on NRT gum should chew slowly for 35 ± 5 minutes, while spacing between each chew
	Inclusion/exclusion criteria	
	Enrolment	
Prior to 9:00 PM	Dinner	
11:00 PM	End of smoking period	

Abbreviations: AE = Adverse event; BMI = Body mass index; CO = Carbon monoxide; CYP2A6 = Cytochrome P450 2A6; mCC = menthol conventional cigarette(s); NRT gum = Nicotine replacement therapy gum; SAE = Serious adverse event; THS 2.2 Menthol= Tobacco Heating System

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

9.3 Investigational Period

9.3.1 Days of Smoking Abstinence (Day 0 and Day 2)

On the days of SA (Day 0 and Day 2) the following assessments will be performed ([Table 6](#) and [Table 7](#)):

Table 6. Time Schedule – Day 0 Washout

Time	Blood sample	Procedures	Additional information
Start of procedure			
		Nicotine abstinence	All day
		Support during nicotine abstinence as required	All day
		AE/SAE recording, concomitant medication	All day
		Randomization	At any time of the day
06:30AM-09:00 AM		Assessment of cough	
Prior to 10:00 AM		Breakfast	
8:00 AM-9:30 AM		CO breath test	
10:00 AM-11:30 AM		Vital signs	At least 5 min in supine position prior to measurement
12:00 PM-1:30 PM		CO breath test	
Prior to 2:30 PM		Lunch	
4:00 PM-5:30 PM		CO breath test	
Afternoon		Snacks	
Prior to 9:00 PM		Dinner	

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

8:00 PM-9:30 PM

CO breath test

Abbreviations: AE = Adverse event; CO = Carbon monoxide; SAE = Serious adverse event.

Table 7. Time Schedule – Day 2 Washout

Time	Blood sample	Procedures	Additional information
Start of procedure			
		Nicotine abstinence	All day
		Support during nicotine abstinence as required	All day
		AE/SAE recording, concomitant medication	All day
	√	Plasma nicotine PK sample	24 hrs +5 min after T ₀ of Day 1 (T ₁₅)
		Assessment of cough	24 hrs minus 5 min after T ₀ of Day 1
Prior to 10:00 AM		Breakfast	
8:00 AM-9:30 AM		CO breath test	
10:00 AM-11:30 AM		Vital signs	At least 5 min in supine position prior to measurement
12:00 PM-1:30 PM		CO breath test	
Prior to 2:30 PM		Lunch	
4:00 PM-5:30 PM		CO breath test	
Afternoon		Snacks	
Prior to 9:00 PM		Dinner	
8:00 PM-9:30 PM		CO breath test	

Abbreviations: CO = Carbon monoxide; PK = Pharmacokinetic; T = Time point.

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

9.3.2 Days of Single Use (Day 1 and Day 3)

On the days of single use (Day 1 and Day 3) the following assessments will be performed ([Table 8](#)):

Table 8. Time Schedule – Day 1 and Day 3 Single Use

Time	Blood sample	Procedures	Additional information
Start of procedure			
		Optional: light snacks prior to first blood draw of the day	
		AE/SAE recording, concomitant medication	All day
		Craving questionnaire (QSU-brief)	For subjects on THS 2.2 Menthol or on mCC: First sample within 15 min prior to T ₀ , 9 assessments thereafter in relation to T ₀ : 15 min, 30 min, 45 min, 60 min, 2 hr, 4 hrs, 6 hrs, 9 hrs, and 12 hrs after T ₀ , with an allowed time window of +10 min each For subjects on NRT gum: First sample within 15 min prior to T ₀ , 9 assessments thereafter in relation to T ₀ : 20 min, 30 min, 45 min, 60 min, 2 hr, 4 hr, 6 hrs, 9 hrs, and 12 hrs after T ₀ , with an allowed time window of +10 min each

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

√	Plasma nicotine sample	<p>For subjects on THS 2.2 Menthol or on mCC: a total of 16 blood samples will be taken for a 24-hour profile (Day 1 and Day 3). The first blood sample will be taken within 15 min prior to T₀ and 15 samples thereafter in relation to T₀: T₁ after 2 min +1 min, T₂ after 4 min +1 min, T₃ after 6 min +1 min, T₄ after 8 min +1 min, T₅ after 10 min +1 min, T₆ after 15 min +2 min, T₇ after 30 min +2 min, T₈ after 45 min +2 min, T₉ after 60 min +3 min, T₁₀ after 2 hrs +5 min, T₁₁ after 4 hrs +5 min, T₁₂ after 6 hrs +5 min, T₁₃ after 9 hrs +5 min, T₁₄ after 12 hrs +5 min, and T₁₅ after 24 hrs +5 min (this sample will be drawn during the day following product use, i.e., wash-out)..</p> <p>For subjects on NRT gum: A total of 16 blood samples will be taken for a 24-hour profile (Day 1 and Day 3). The first blood sample will be taken within 15 min prior to T₀ and 15 samples thereafter in relation to T₀: T₁ after 10 min + 1 min, T₂ after 20 min + 1 min, T₃ after 25 min + 1 min, T₄ after 30 min + 1 min, T₅ after 35 min + 1 min, T₆ after 40 min + 1 min, T₇ after 45 min + 1 min, T₈ after 60 min + 3 min, T₉ after 2 hrs + 5 min, T₁₀ after 3 hrs + 5 min, T₁₁ after 4 hrs + 5 min, T₁₂ after 6 hrs + 5 min, T₁₃ after 9 hrs + 5 min, T₁₄ after 12 hrs + 5 min, and T₁₅ after 24 hrs +5 min (this sample will be drawn during the day following product use, i.e., wash-out).</p>
√	COHb blood sampling	Five blood samples to be taken, first sample within 15 min prior to T ₀ , then after 15 min +2 min, 60 min +3 min, 4 hrs +5 min, 12 hrs +5 min
	CO breath test	First test to be done within 15 min prior to T ₀

	Assessment of cough	Has to be done prior to product use
6:00 AM-9:00 AM	Start of single product use	
Prior to 10:00 AM	Breakfast	
10:00 AM-11:30 AM	Vital signs	At least 5 min in supine position prior to measurement
10:00 AM-11:30 AM	ECG	At least 10 min in supine position prior to recording
12:00 PM-1:30 PM	CO breath test	
Prior to 2:30 PM	Lunch	
4:00 PM-5:30 PM	CO breath test	
Afternoon	Snacks	
Prior to 9:00 PM	Dinner	
8:00 PM-9:30 PM	CO breath test	
8:00 PM-11:00 PM	Product evaluation questionnaire (MCEQ; only in sequence 1 and 2)	
	Collection of used Menthol Tobacco Sticks and mCC butts and NRT gum	After the product use

Abbreviations: AE = Adverse event; CO = Carbon monoxide; COHb = Carboxyhemoglobin; ECG = Electrocardiogram; mCC = menthol conventional cigarette(s); MCEQ = Modified Cigarette Evaluation Questionnaire; NRT gum = nicotine replacement therapy gum; QSU-brief = Questionnaire of Smoking Urges brief; SAE = Serious adverse event; T = Time point; THS 2.2 Menthol = Tobacco Heating System 2.2 Menthol

9.4 Day of Discharge

The following assessments will be conducted prior to the time of Discharge on Day 4 (or after a subject is prematurely withdrawn from the study) ([Table 9](#)):

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

Table 9. Time Schedule – Day 4 Discharge

Time	Blood sample	Procedures	Additional information
Start of procedure			
		AE/SAE recording, concomitant medication	All day
	√	Plasma nicotine: PK sample	24 hrs +5 min since T ₀ of Day 3 (T ₁₅)
		Assessment of cough	24 hrs minus 5 min since T ₀ of Day 3
	√	Clinical laboratory parameters (hematology, urine analysis, clinical chemistry)	Has to be done in at least 10 hour fasting condition
		Spirometry	Has to be done prior to smoking
prior to 10:00 AM		Breakfast	
Prior to discharge		CO breath test	
Prior to discharge		Urine pregnancy test in all female	
Prior to discharge		Vital signs	At least 5 min in supine position prior to measurement
Prior to discharge		Physical examination, weight and calculated BMI	
Prior to discharge		Advice on risk of smoking and debriefing	
Time of discharge			

Abbreviations: AE = Adverse event; BMI = Body mass index; CO = Carbon monoxide; PK = Pharmacokinetic; SAE = Serious adverse event; T = Time point.

Smoking will be allowed after spirometry has been conducted.

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

9.5 Safety Follow-up Period

All subjects participating in the product trial on Day -1 and are not enrolled into the study will enter a 7-day safety follow-up period.

After the time of Discharge at Day 4 (or if prematurely withdrawn from the study), subjects will enter a 7-day safety follow-up period.

During the 7-day safety follow-up period, there will be spontaneous reporting by the subject of new AEs and new SAEs. Any ongoing AEs/SAEs will be actively followed up by the site.

Any AEs or SAEs that are ongoing at the end of the 7-day safety follow-up period will be handled as described in [Section 8](#).

9.6 Early Termination Procedures

The Day of Discharge assessments will be performed as early termination procedures (see [Section 9.4](#)). Early termination procedures will be the same as those described in the Day of Discharge.

10 QUALITY CONTROL AND QUALITY ASSURANCE

10.1 Monitoring

The Clinical Research Associate (CRA) (“Monitor”) will be responsible for the monitoring of the study. Monitoring will be performed according to CRO’s SOPs and as per the agreed monitoring plan with the Sponsor.

The Principal Investigator/head of the investigational site shall permit the Monitor to review study data as frequently as considered necessary to ensure that data are being recorded in an adequate manner and that protocol adherence is satisfactory.

The Investigator shall access medical records for the Monitor in order that entries in the eCRFs may be verified. The Investigator, as part of their responsibilities, is expected to ensure that the study adheres to GCP requirements.

An Investigator’s meeting will be held prior to the site initiation visit. During this meeting, the general training of the study procedures and specific training on selected procedures will be done and documented.

Subsequent to the Investigator’s meeting, and before the first subject is screened into the study, site initiation visit will be conducted by the Monitor and, if necessary, with Sponsor or its authorized representative. The purpose of the site initiation visit will be detailed in the monitoring plan.

Communication by telephone, mail, and e-mail may be used as needed to supplement site visits. The Investigator and study personnel will cooperate with the Monitor, provide all appropriate documentation, and will be available to discuss the study.

The Monitor and the Sponsor’s personnel will be available between visits, should the Investigator or other study collaborator at the sites need information and advice.

Site visits will be made at regular intervals during the study. The frequency of the monitoring visits will be defined in the monitoring plan agreed with the Sponsor.

The Investigator, or a designated member of the Investigator’s study collaborator, must be available during the monitoring visit to review the data and resolve any queries, and to allow direct access to the subject’s records for source data verification.

10.2 Training of Study Collaborator

A formal meeting (Investigator's meeting) will be conducted prior to site initiation. During this meeting, the Sponsor or its authorized representative will discuss the requirements of the clinical study protocol and related documents and will also provide training in the relevant systems and other study-specific procedures. The activities of the Investigator's meeting will be described in the monitoring plan.

In addition to the Investigator meeting, the Investigator will ensure that appropriate training relevant to the study is provided to all study collaborator involved in the study, and that any new information relevant to the performance of this study is forwarded to the study collaborator involved in a timely manner. The record of all individuals involved in the study will be maintained in the Site Investigator File.

10.3 Audits and Inspections

Good Clinical Practice regulations require that there are independent inspections of clinical program activities. Such inspections may be performed at any time before, during and/or after the study.

Authorized representatives of the Sponsor, regulatory agencies and/or an IRB may perform audits or inspections, including source data verification. The purpose of an audit or inspection is to systematically and independently examine all study-related activities and documents to determine whether these activities were conducted, and data were recorded, analyzed, and accurately reported according to the protocol, ICH/GCP guidelines, and any applicable regulatory requirements. The Principal Investigator/head of investigational site will contact the Sponsor or the authorized representative immediately, if contacted by a regulatory agency about an inspection at their site.

The Investigator and study collaborator are responsible for maintaining a comprehensive and accurate filing system of all study-related documentation that will be suitable for inspection at any time by the Sponsor, its authorized representative, and/or regulatory agencies. In signing this protocol, the Investigator understands and agrees to provide access to the necessary documentation and files.

11 DATA MANAGEMENT ACTIVITIES

All Data Management Activities will be described in detail in the Data Management Plan and documents specified therein.

11.1 Data Capture

11.1.1 Case Report Forms and Study Records

With the exception of the subject reported outcome data, all results from the clinical assessments will be recorded in the Source Documents by the Investigator and then captured in the eCRFs at the study site. The subject questionnaires and the VAS will be entered by the subject directly in the ePRO device. Trained study personnel will be responsible for capturing the data from the observations, tests, and assessments specified in the protocol in the Source Documents and then transferring the data into the eCRF according to the eCRF Completion Guidelines.

The Principal Investigator has ultimate responsibility for the collection and reporting of all data related to the clinical study and ensuring that the data are accurate, authentic/original, legible, timely (contemporaneous), enduring, and available when required. The eCRF must be signed by the Investigator to attest that the data contained in the eCRF are true and accurate. Any corrections made to Source Documents must be clearly recorded, without obscuring the original values and be accompanied by the date of change, reason for change, and identification of the person making the change. The eCRF for each subject will be checked against the Source Documents at the study site by the CRA. Instances of missing or unclear data will be discussed with the Investigator for resolution. An eCRF will be generated for all subjects that sign the informed consent.

11.1.2 Protocol Deviations

All protocol deviations will be entered into the Clinical Trial Management System (CTMS) or other approved format.

Information from the Source Documents will represent the primary source of protocol deviations. Information following site monitoring and other manual reviews will be documented in the site visit reports, follow-up letters, audit documentation, or other manual review and will be recorded and tracked in the CTMS or other approved format. Telecommunications and other verbal communications regarding deviations will be considered and handled as important communication, documented and tracked as protocol deviations, as necessary.

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

Individual entries for protocol deviations that are recorded in the CTMS, or other approved format, following site monitoring and other manual reviews will be reviewed against the individual data points in the eCRF database but will not be formally reconciled with the eCRF database (e.g., their description or occurrence date). The overall procedure for managing protocol deviations are described in the SOPs of the CRO Data Management Team. All deviations will be reviewed periodically, as determined at study start, to identify trends to improve monitoring and/or potential impact on the statistical analysis.

11.2 Data Handling

All study data will be managed by the Data Management Team at the CRO. The overall procedures for quality assurance of clinical study data are described in the SOPs of the CRO Data Management Team. The Data Management Team at CRO will prepare a Data Management Plan (DMP), to be reviewed and approved by the Sponsor, prior to the start of the study. This document will describe, in detail, the Data Management related procedures and processes.

All data of all subjects enrolled and screening failures who experience an AE during the study (from time of informed consent to end of the safety follow-up period) will be captured and stored in the study database.

All data collected during the study is property of the Sponsor irrespective of the location of the database and the Data Management CRO.

11.2.1 Data Validation

The data will be validated as defined in the DMP and Data Validation Specifications. Discrepancies will be reported as defined in DMP and Data Validation Plan.

Data queries will be raised for discrepant or missing data. All changes to data will be captured in the database with a comprehensive audit trail.

11.2.2 Coding

Adverse events, medical/surgical history, and prior/concomitant medication will be classified according to the terminology of the latest version of the following Dictionaries, at time of coding the first entry:

Medical history:	Medical Dictionary for Regulatory Activities (MedDRA®)
Adverse events:	MedDRA®
Medications:	WHO Drug Dictionary Enhanced and Anatomical Therapeutic and Chemical classification system

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

THS 2.2Menthol device C54451/Medical_Device_Problem_Codes_FDA_CDRH
issues and/or
malfunctions:

11.2.3 Database Lock

When all outstanding Data Management issues have been resolved and all validation, quality review, and cleaning activities are complete, the database or selected data is/are declared soft locked. Access to change data in the soft-locked database or to change selected data at this time is limited.

After the data is reviewed by the Sponsor, resolution of all raised queries and QC of the changed data, database, or selected data upon Sponsor approval as applicable, is declared locked.

Any changes to the database after that time can only be made by written agreement between the Sponsor and the Data Management and Statistical Team at the CRO. Any of those changes must be documented in the database log file.

After study completion, the study database will be transferred to the Sponsor in the format specified in the DMP in Clinical Data Interchange Standards Consortium's Study Data Tabulation Model Data Structure Specifications.

12 PLANNED STATISTICAL METHODS

12.1 General Considerations

An incomplete block design will be adopted in this study with every subject being exposed to 2 out of the 3 study products (mCC, THS 2.2 Menthol, and NRT gum) to allow comparisons between THS 2.2 Menthol and mCC in Group-1 (sequence 1 and 2), and between THS 2.2 Menthol and NRT gum in Group-2 (sequence 3 and 4).

Full details of the statistical analysis will be given in a SAP. Any changes to the planned statistical methods will be documented in the Clinical Study Report. The statistical evaluation will be performed using SAS[®], version 9.2 or later.

12.1.1 Stratification Criteria

For analysis, the following stratification criteria will be used:

- Sex (male and female)
- mCC nicotine level at Admission (ISO nicotine levels ≤ 0.6 mg and 0.6 to 1 mg)

In addition, for the safety data, the analysis will be stratified by sequence and by study periods (Screening, product test, product exposure, and safety follow-up period).

12.1.2 Definitions for Statistical Data Analysis

Unless otherwise stated, for the purposes of statistical analyses, baseline is defined as the last available time point prior to T₀ on Day 1, from 6:00 AM to 9:00 AM.

12.1.3 Descriptive Statistics

All data will be presented in listings, ordered by sequence and subject, unless otherwise specified.

Descriptive statistics for continuous variables (number of subjects [n], number and percentage of subjects with missing data, mean, standard deviation, median, first and third quartiles, minimum and maximum for continuous data, and the n and absolute and relative [%] frequency for categorical data) will be presented by exposure and overall at each time point, where applicable.

Descriptive statistics for PK parameters will also include the geometric mean and coefficient of variation (CV).

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

Baseline, admission and screening data (i.e., anything prior to product exposure) will be summarized by sequence and overall where appropriate

12.1.4 Handling of Missing Values and of Values outside the Detection Limits

In general, missing data will not be imputed in this study, due to the nature of the measurements and the short periods of exposure and use. However, for questionnaire data total scores and domain or subscale scores may use a certain degree of imputation by averaging across individual item scores. Further details will be provided in the SAP.

Values below the lower limit of quantification (LLOQ) will be imputed using $0.5 \times \text{LLOQ}$. For values above the upper limit of quantification (ULOQ), i.e., preceded by a ">", for example ">xx", the numerical xx will be used for calculation and reporting in summary tables. The number of values below LLOQ or above ULOQ will be presented in each summary table.

12.1.5 Significance Level for Inferential Analysis

For all endpoints, unless otherwise stated, statistical tests will be two-sided and conducted at the 5% significance level and all quoted confidence intervals will be two-sided 95% CIs.

No formal adjustment of the test-wise alpha level for multiple testing is necessary, as no claim will be made based on the outcome of the individual tests.

12.2 Determination of Sample Size and Power Consideration

A total of 62 subjects will be randomized. This sample size is the sum of the sample size requirements for the THS 2.2 Menthol: mCC comparison and the THS 2.2 Menthol:NRT gum comparison (as described in detail below).

The estimates for the within-subject CV for nicotine C_{\max} (36%) and $\text{AUC}_{(0-\text{last})}$ (21%) are based on the data collected in the ZRHX-PK-02 clinical study ([ZRHX-PK-02, 2012](#)) comparing the nicotine PK profiles of Tobacco Heating System 2.1, the predecessor of THS 2.2 Menthol (regular tobacco stick) and CC. In the absence of data comparing THS and NRT gum, the same CVs were assumed for the calculation of the sample size related to the THS 2.2 Menthol:NRT gum comparison.

Anticipated mean C_{\max} and $\text{AUC}_{(0-\text{last})}$ for THS 2.2 Menthol and mCC were based on data from the ZRHX-PK-02 study ([ZRHX-PK-02, 2012](#)). Anticipated mean C_{\max} and $\text{AUC}_{(0-\text{last})}$ for NRT gum were based on data reported by [Dautzenberg et al., 2007](#).

The sample sizes of this study assume a drop-out rate no larger than 10%.

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

Sample size calculations were conducted using SAS[®] version 9.2 for the 95% CI of the mean differences between paired observations (proc power onesamplemeans) in the natural log scale (Senn, 2002). The SAS[®] implementation of the method published by Beal, 1989 was adopted to estimate the probability of obtaining at most the target 95% CI of $\pm 20\%$.

THS 2.2 Menthol:mCC comparison

A total of 44 subjects are needed to estimate the mean C_{\max} parameter ratio between THS 2.2 Menthol and mCC with a 90% probability of obtaining a margin of error (95% CI) of at most $\pm 20\%$, assuming that THS 2.2 Menthol have a nicotine C_{\max} similar to mCC (ratio equal to 1.00). This sample size is sufficient to provide 90% probability of obtaining a margin of error of at most $\pm 20\%$ for the $AUC_{(0-\text{last})}$ ratio between THS 2.2 Menthol and mCC, assuming a similar extent of nicotine absorption for the 2 products (ratio equal to 1.00).

THS 2.2 Menthol:NRT gum comparison

A total of 18 subjects are needed to estimate the mean $AUC_{(0-\text{last})}$ parameter ratio between THS 2.2 Menthol and NRT gum with a precision allowing for the lower bound of the 95% CI exceeding 1.00, with 90% power, assuming a geometrical ratio between THS 2.2 Menthol and NRT gum of 1.28. This sample size is sufficient to provide 90% power of obtaining a lower bound of the 95% CI of C_{\max} ratio between THS 2.2 Menthol and NRT gum, exceeding 1.00, assuming a geometrical mean ratio between THS 2.2 Menthol and NRT gum of 2.00.

12.3 Analysis Populations

All analyses will be based on actual product exposure. All endpoints (other than safety) will be analyzed using the PK analysis sets. Safety will be analyzed using the safety population.

12.3.1 PK Populations

The analysis populations for the PK endpoints are composed of two analysis sets to allow the comparison between THS 2.2 Menthol and NRT gum separately from the comparison between THS 2.2 Menthol and mCC.

The PK populations consist of all the randomized subjects who give informed consent, completed at least one of the single use Day 1 or Day 3, and for whom at least one PK parameter can be derived. Only subjects without major protocol deviations (to be defined in the SAP) will be included in the PK analysis sets.

12.3.2 Safety Population

The safety population consists of all the subjects who give informed consent and have at least one exposure to THS 2.2 Menthol (including the product test at Admission Day).

12.4 Demographics and Baseline Characteristics

Demographic and other baseline characteristics will be reported for the PK and safety populations. Appropriate summary statistics will be provided, as described in [Section 12.1.3](#).

12.5 Primary Endpoints

12.5.1 Primary Endpoint Analysis Variables

Nicotine PK parameters will be derived from plasma nicotine versus time data using a non-compartmental technique. In particular:

C_{\max}	Maximum observed plasma concentration. C_{\max} will be reported as long as there is at least one quantifiable concentration post-exposure
$AUC_{(0-\text{last})}$	Area under the plasma concentration-time curve from start of product use to the time of the last quantifiable concentration (linear trapezoidal method).

12.5.2 Baseline Comparability

Not applicable.

12.5.3 Descriptive Analysis

Primary endpoints will be summarized as described in [Section 12.1.3](#).

An analysis of variance (ANOVA) will be conducted on $AUC_{(0-\text{last})}$ and C_{\max} endpoints in the natural logarithmic scale. The model will include terms for sequence, subjects within sequence, period and product exposure as fixed effect factors. The results of this analysis for each of $AUC_{(0-\text{last})}$ and C_{\max} are presented in terms of adjusted geometric least square means and 95% CIs for the THS 2.2 Menthol:mCC and THS 2.2 Menthol:NRT gum ratios.

This approach is consistent with the guidelines in the European Medicines Agency's guidelines for bioequivalence investigations ([EMA, 2008](#)) and FDA's Center for Drug Evaluation and Research ([FDA, 2001](#)). Carry-over effect will not be tested, as it cannot be statistically distinguished from the interaction between treatment and period in a 2x2 crossover design ([ICH E9, 1998](#)).

A sensitivity analysis will be conducted should there be 20% or more missing PK parameter values, by repeating the above analyses using mixed effects ANOVA model in the natural log scale, with a restricted maximum likelihood method to estimate mean differences and variances as suggested by FDA ([FDA, 2001](#)). Subjects within sequence will be used as Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

random effects and fixed effects are period, sequence, and product exposure. To evaluate the sensitivity to the distributional assumptions, point and interval estimates will also be estimated by means of the percentile bootstrap technique, using 2000 bootstrap samples which preserve the number of subjects per sequence.

12.5.4 Confirmatory Analyses

Given that the objective of this study is to determine the point estimate and precision of the ratio of THS 2.2 Menthol:mCC for C_{\max} and $AUC_{(0-\text{last})}$, there is no statistical hypothesis to be tested.

12.6 Secondary Endpoints

12.6.1 Secondary Endpoint Analysis Variables

Nicotine PK parameters will be derived as follows:

t_{\max} Time of maximum observed plasma concentration. t_{\max} will be reported as long as there is at least one quantifiable concentration post-exposure

$AUC_{(0-t^*)}$ Area under the plasma concentration-time curve from start of product use to the subject-specific time of maximum nicotine concentration following mCC or NRT gum (linear trapezoidal method)

$AUC_{(0-\infty)}$ Area under the plasma concentration-time curve from start of product use extrapolated to infinite time, according to:

$$AUC_{0-\infty} = AUC_{0-\text{last}} + \left(\frac{C_{\text{last}}}{\lambda_z} \right)$$

Where C_{last} is the last quantifiable concentration and λ_z is the terminal elimination rate constant

λ_z Terminal elimination rate constant, estimated by linear regression analysis of the natural log-transformed concentration-time data

$t_{1/2}$ Terminal elimination half-life, derived as $\ln(2)/\lambda_z$

More details on PK parameter derivations will be provided in the SAP.

The subjective effects of using THS 2.2 Menthol as compared to mCC and NRT gum will be evaluated by analyzing domain scores of the QSU-brief and MCEQ. Full details of questionnaire domain scores derivation will be provided in the SAP.

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

12.6.2 Baseline Comparability

Not applicable.

12.6.3 Descriptive Analysis

In general, secondary endpoints will be summarized using the approach described in [Section 12.1.3](#).

The following analyses will be conducted in both Group-1 and Group-2 PK analysis sets:

- The $AUC_{(0-\infty)}$, $AUC_{(0-t^*)}$, and $t_{1/2}$ will be analyzed using the same approach adopted for the primary endpoints. No statistical analysis will be performed on the elimination rate constant λ_z . Data will be summarized and presented together with 95% CI.
- The Hodges-Lehmann 95% CI estimates for the median t_{max} differences will be presented ([Lingling, 2008](#)).
- The use of random effects model is suggested in the context of the analysis of subjective effects of smoking ([Shiffman et al, 2004](#)). Mixed effects ANOVA with the same model terms as planned for the sensitivity analysis of the primary endpoints will be adopted for the analysis of QSU-brief, including all of the different assessment time points as repeated measurements. The analysis will not be adjusted for the assessment prior to T_0 ([Fleiss et al, 1985](#)).
- Levels of exhaled CO and of blood COHb will be summarized by means of descriptive statistics reported by exposure. Analysis of COHb levels will be conducted using a mixed model for repeated measures, the same approach as for the QSU-brief.

The following analyses will be conducted only in Group-1 PK analysis set:

- Mixed effects ANOVA with the same model terms as planned for the sensitivity analysis of the primary endpoints will be adopted for the analysis of the MCEQ domain scores.

The following analyses will be conducted in only Group-2 PK analysis set:

- To test if the time to the maximum nicotine concentration in THS 2.2 Menthol is shorter than in NRT gum the following hypothesis will be evaluated:

$$H_0: X_{THS} - X_{NRT} = 0 \quad H_A: X_{THS} - X_{NRT} < 0$$
 where X_{THS} and X_{NRT} are the median values of the THS 2.2 Menthol and NRT gum, respectively. The t_{max} will be analyzed on the original scale using the Wilcoxon

Signed-Rank Test with a type I error $\alpha = 0.025$ (one-sided test), as values are ordinal/discrete, and the assumption of normality may be questionable.

- To determine if the rate and the amount of nicotine absorbed of the THS 2.2 Menthol is higher relative to NRT gum the following hypothesis will be tested for both C_{\max} and $AUC_{(0-\text{last})}$ parameters:

$$H_0: X_{THS}/X_{NRT} = 1.0 \quad H_A: X_{THS}/X_{NRT} > 1.0$$

where X_{THS} and X_{NRT} are the adjusted geometrical means of THS 2.2 Menthol and NRT gum, respectively. H_0 is rejected with a type I error $\alpha = 0.025$ (one-sided test), if the lower bound of the 95% CI for the X_{THS}/X_{NRT} ratio is higher than 1.0.

12.6.4 Confirmatory Analyses

Not applicable.

12.6.5 Safety Endpoints

In general, all safety data will be listed and tabulated on the safety population by sequence, using the approach described in [Section 12.1.3](#). Safety variables collected during exposure periods will also be reported by product exposure.

AE data will serve as the primary assessment of safety. Other safety variables monitored in this study include: respiratory symptoms (cough assessment, VAS, and Likert scales); vital signs (systolic and diastolic blood pressure, pulse rate, respiratory rate); spirometry; ECG data; clinical chemistry, hematology, and urine analysis safety panel; physical examination; and concomitant medications.

The number and percentage of subjects with AEs and SAEs will be tabulated by system organ class and preferred term. Summaries will also be presented for AEs leading to withdrawal, AEs leading to Death, AEs by relatedness to product exposure, AEs by severity, and laboratory AEs. Tabulations will be performed for both the number of subjects experiencing an event and the number of events. The number and percentage of device events and of subjects with device events will be tabulated by sequence.

The number and percentage of subjects with clinical findings will be tabulated by sequence for laboratory parameters. Shift tables showing change from baseline of clinical findings will be provided for: ECGs, physical examinations, and laboratory parameters (both shifts in normal ranges and toxicity grades). Descriptive statistics will be summarized by visit and change from baseline for laboratory parameters, ECG, respiratory symptoms, and vital signs.

12.7 Exploratory Analyses

There are no planned exploratory analyses.

12.8 Interim Analysis

There are no planned interim analyses.

13 ADMINISTRATIVE CONSIDERATIONS

13.1 Principal Investigators and Study Administrative Structure

13.1.1 Principal Investigator

Principal Investigator: Fumimasa Nobuoka M.D.	Ageo Medical Clinic 3133 Haraichi, Ageo City Saitama 362-0021, Japan Tel:+81-48-720-2730
---	--

13.1.2 Sponsor

Sponsor:	Philip Morris Products S.A Quai Jeanrenaud 5 2000 Neuchâtel Switzerland. Phone: +41 (58) 242 2111 Fax: +41 (58) 242 2811
Christelle Haziza, PhD Manager Clinical Science	Phone: +41 (58) 242 2625 Mobile: +41 (79) 336 1781 E-mail: Christelle.Haziza@pmi.com
Nicola Lama, PhD Staff Scientist, Statistics	Phone: +41 (58) 242 2694 Mobile: +41 (79) 700 9374 Email: Nicola.Lama@pmi.com
Tamara Koval, MD Medical Safety Officer	Phone: +41 (58) 2422112 Mobile: +41 (79) 484 90 68 E-mail: Tamara.Koval@pmi.com
Lars Olsen Medical Writer	Phone: +41 (58) 242 2532 Mobile: +41 (79) 485 0371 Email: Lars.Olsen@pmi.com
Muriel Benzimra, MSc Clinical Study Manager	Phone: +41 (58) 222 2638 Mobile: +41 (79) 8159 628 E-mail: Muriel.Benzimra@pmi.com

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

13.1.3 Other Responsibilities

(b) (4)

(b) (4) is the local Contract Research Organization designated by PMI to manage and monitor the study: all duties and responsibilities transferred to (b)(4) by PMI will be defined in the agreement signed between the two parties

(b) (4)

Tel: (b) (4) 4 / Fax: (b) (4)

Responsible person (Clinical Pharmacology): (b) (4)

(b) (4)

Tel: (b) (4) / Fax: (b) (4)

Any SAEs or pregnancies will be handled by:

UBC Safety
16, Chemin des Coquelicots
1214 Vernier/Geneva
Switzerland
Phone: +41 (22) 596 4444
Fax: +41 (22) 596 4446
E-mail: EUsafety@unitedbiosource.com

Details of the laboratories conducting the clinical safety laboratory services, and biopharmaceutical analyses are shown in [Appendix 2](#).

13.2 Subject Confidentiality

All information obtained during the conduct of the study with respect to the subjects' state of health will be regarded as confidential. A statement to this effect will be written in the information provided to the subject. An agreement to disclose any such information will be obtained from the subject in writing and signed by the subject, in compliance with all local and national data protection and privacy legislation.

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

The anonymity of subjects participating in this study will be maintained. Subjects will be identifiable by the Sponsor (or Sponsor's authorized representative) on eCRFs and other documents by their subject (or randomization) number/code, sex and date of birth, but **not** by name, initial, or any other details relating to identifiable person (e.g. address, health insurance ID card, medical chart number, etc.). The assignment of a subject number/code for subject identification will be based on the appropriate data protection rules.

Any documents that allow full identification of the subject (e.g. the subject's signed ICF) must be maintained in confidence by the Investigator. If any document relating to this study shows a subject's name or any other details relating to an identifiable person (e.g. address, health insurance ID card, medical chart number, etc), the name or other identifiable details must be obscured before a copy of that document is supplied to the Sponsor or the Sponsor's authorized representative.

13.3 Access to Source Documentation

Subjects will be informed that, during the course of the clinical study, the Sponsor, any authorized representatives of the Sponsor, IRB, or regulatory authorities may inspect their medical records to verify the information collected, and ensure that all personal information made available for inspection is handled in the strictest confidence and in accordance with national and local data protection and privacy legislation.

The Investigator/head of the investigational site and all study site staff involved with the study must permit direct access to source data/documents for study related monitoring, audits, IRB review, and regulatory inspection(s).

13.4 Record Retention

All records of data, source data and source documents (original records or certified copies), in any form (including, but not limited to, written, electronic, magnetic, optical records and scans, X-rays, and ECGs) that describe or record the methods, conduct, and/or results of the study, the factors affecting the study, and the actions taken will be maintained by the Principal Investigator/head of investigational site for the study, as required by ICH GCP and any other applicable local or national regulations. For X-rays at least the radiologist's assessment is required as source documentation. If the actual image is available it can be stored on CD as well.

Essential study documents/records, which individually and collectively permit evaluation of the conduct of a study and the quality of the data produced, are described in Article 41 of Ministerial Ordinance on GCP (Ordinance of the Ministry of Health and Welfare No. 28 of March 27, 1997 (as last amended by the Ordinance of Ministry of Health, Labour and Welfare No. 161 of December 28, 2012). Essential documents must be retained by the Principal Investigator/head of investigational site for a minimum of:

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

- At least 15 years after completion or discontinuation of the study, or
- At least 2 years depending on, for example, the circumstances, or
- After formal discontinuation of clinical development of the IP.

These documents should be retained for a longer period, however, if required by the applicable regulatory requirements or by an agreement with the Sponsor.

Examples of essential records/documents include, but are not limited to:

- Signed informed consent documents for all subjects and Master ICF.
- Subject identification code list, Screening Log and Enrolment Log (if applicable).
- Record of all communications between the Principal Investigator and the IRB, composition of the IRB.
- Record of all communications/contact between the Investigator, Sponsor, and its authorized representatives.
- List of sub-Investigators and other appropriately qualified persons to whom the head of the investigational site has delegated significant study-related duties, together with their roles in the study.
- Investigator Logs.
- eCRFs, study specific questionnaires (and associated data/scoring), subject diaries.
- Adverse event reports and details of follow-up investigations, details of concomitant medication.
- All other Source Documents (e.g., chest X-rays, ECGs, consultation reports, physical examination, laboratory records) or any electronically captured study source data.
- Clinical laboratory reports, laboratory normal ranges.
- Original medical/hospital records, if applicable (the medical files of study subjects must be retained in accordance with local legislation and in accordance with the maximum period of time permitted by the hospital or study site).
- Record of any body fluids or tissue samples collected and retained.
- Device Issue Log, IP Accountability Logs, dispensing records.
- Information regarding subjects' discontinuation and any follow-up.

It is the responsibility of the Sponsor to inform the Principal Investigator/head of investigational site as to when these documents no longer need to be retained.

The Principal Investigator/head of investigational site must take measures to prevent accidental or premature destruction of these documents.

If the head of the investigational site wishes to assign the study records to another party or move them to another location, the Sponsor must be notified in advance.

The head of the investigational site must obtain written approval from the Sponsor before destruction of any records. Normally, these records will be held in the Investigator's archives. If the head of the investigational site is unable to meet this obligation, they must ask the Sponsor for permission to make alternative arrangements. Details of these arrangements must be documented.

The Sponsor or Sponsor's authorized representative will maintain documentation relating to the study as long as the IP is on the market, and/or for 15 years after the CSR has been finalized.

13.5 Clinical Study Report

The Sponsor must ensure that a CSR for this study is prepared regardless of whether the study is completed or prematurely terminated.

The CSR will be written based on standards of the ICH Guideline for the Structure and Content of Clinical Study Reports. In certain circumstances, an abbreviated CSR may be acceptable. Submission of the CSR to the IRB will be complied with as requested by local requirements.

The results of the additional variables for analysis will be presented in reports separate from the study CSR.

13.6 Financial Disclosure

Investigators are required to provide financial disclosure information to the Sponsor. In addition, the Investigators must provide to the Sponsor a commitment to promptly update this information if any relevant changes occur during the course of the investigation and for 1 year following the completion of the study.

13.7 Publication and Disclosure Policy

This document contains data, information, and trade secrets that are confidential and proprietary to the Sponsor. This document is being provided solely for the purpose of evaluation and/or conducting this clinical study for the Sponsor. Disclosure of the content of this document is allowed only to study personnel, IRB, or duly authorized representatives of regulatory agencies for this purpose under the condition that confidentiality is maintained. The contents of this document may not be used in any other clinical study, disclosed to any Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

other person or entity without the prior written permission of the Sponsor. The foregoing shall not apply to disclosure required by any regulations; however, prompt notice will be given to the Sponsor prior to any such disclosure.

The Sponsor plans to disclose details of the study protocol on a web-based, publicly available, clinical trial register database (e.g., ClinicalTrials.gov).

13.8 Insurance

The Sponsor is responsible for AEs and health damage of the subjects that are associated with the THS 2.2 Menthol product or with study procedures which are used during the study, except for AEs and health damage of the subjects caused by a negligent or an intentional misconduct and/or significant deviation to the protocol of the Investigator or the clinical trial site or the subjects. The Sponsor has taken out insurance to cover any bodily injury and property damage caused by the operations carried out by the insured.

14 REFERENCE LIST

Beal, 1989

Beal SL. Sample Size Determination for Confidence Intervals on the Population Means and on the Difference between Two Population Means. 1989; Biometrics, 45, 969–977.

Benowitz et al. 2002

SNRT subcommittee on biochemical verification. Biochemical verification of tobacco use and cessation. Nicotine Tob Res. 2002; 4 (2): 149-159.

Benowitz et al, 2006

Benowitz NL, Lessov-Schlaggar CN, Swan GE, Jacob P 3rd. Female sex and oral contraceptive use accelerate nicotine metabolism. Clin Pharmacol Ther. 2006;79(5):480-488.

Benowitz et al, 2009

Benowitz NL, Hukkanen J, Jacob P. Nicotine chemistry, metabolism, kinetics and biomarkers. Handb Exp Pharmacol. 2009;192:29-60.

Cappelleri et al., 2007

Cappelleri JC, Bushmakina AG, Baker CL, Merikle E, Olufade AO, Gilbert DG. Confirmatory factor analyses and reliability of the modified cigarette evaluation questionnaire. Addict Behav. 2007 May; 32 (5): 912-23.

Connolly et al., 2011

Connolly GN, Behm I, Osaki Y, Wayne GF. The impact of menthol cigarettes on smoking initiation among non-smoking young females in Japan. Int J Environ Res Public Health. 2011; 8 (1): 1-14.

Cox et al. 2001

Cox LS, Tiffany ST, Christen AG. Evaluation of the brief questionnaire of smoking urges (QSU-brief) in laboratory and clinical settings. Nicotine Tob Res. 2001; 3 (1): 7-16.

Dautzenberg et al., 2007

Bertrand Dautzenberg, Mitchell Nides, Jean-Luc Kienzler and Anne Callens. Pharmacokinetics, safety and efficacy from randomized controlled trials of 1 and 2 mg

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

nicotine bitartrate lozenges (Nicotinell®). BMC Clinical Pharmacology 2007, 7:11 doi:10.1186/1472-6904-7-11. Available at <http://www.biomedcentral.com/content/pdf/1472-6904-7-11.pdf> (accessed on 01 February 2013)

Declaration of Helsinki (2008)

Declaration of Helsinki. World Medical Association. Available from: <http://www.wma.net/en/30publications/10policies/b3/index.html> (accessed on 21 March 2012).

EMA, 2008

European Medicines Agency, Committee for Medicinal Products for Human Use (CHMP). Guideline on the investigation of bioequivalence. 2008. CHMP/EWP/QWP/1401/98 Rev. 1. Available at http://www.ema.europa.eu/docs/en_GB/document_library/Scientific_guideline/-2009/09/WC500003011.pdf

Fagerström et al., 2012

Fagerström K, Russ C, Yu C, Yunis C and Foulds J. The Fagerström Test for Nicotine Dependence as a Predictor of Smoking Abstinence: A Pooled Analysis of Varenicline Clinical Trial. Nicotine & Tobacco Research. 2012; first published online March 30, 2012.

Fagerström and Schneider, 1989

Fagerström KO, Schneider NG. Measuring nicotine dependence: a review of the Fagerström Tolerance Questionnaire. Journal of Behavioral Medicine. 1989; 12: 159-181.

FDA, 2001

United States Food and Drug Administration, Center for Drug Evaluation and Research. Statistical Approaches to Establishing Bioequivalence. January 2001.

Available from <http://www.fda.gov/downloads/Drugs/.../Guidances/ucm070244.pdf>

FDA, 2002

United States Food and Drug Administration. Guidance for Industry, Food-Effect Bioavailability and Fed Bioequivalence Studies, 2002.

Available from <http://www.fda.gov/cder/guidance/index.htm> (accessed on 26 Sep 2012).

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

FDA, 2011

United States Food and Drug Administration, Harmful and Potentially Harmful Constituents in Tobacco Products and Tobacco Smoke; Request for Comments (76 FR 50226, Aug. 12, 2011).

FDA, 2012a

United States Food and Drug Administration, Center for Tobacco Products. Modified Risk Tobacco Product Applications: Draft Guidance for Industry, March 2012. Available from <http://www.fda.gov/TobaccoProducts/GuidanceComplianceRegulatoryInformation/default.htm>.

FDA, 2012b

United States Food and Drug Administration. Implementation Specifications, reporting medical device problems, C54451/Medical Device Problem Codes, FDA, CDRH. Available from <http://www.fda.gov/MedicalDevices/Safety/ReportaProblem/EventProblemCodes/ucm135741.htm#> (accessed on 04 October 2012).

FDA, 2012c

United States Food and Drug Administration, Center for Tobacco Products. Modified Risk Tobacco Product Applications: Draft Guidance for Industry, March 2012.

Fleiss et al, 1985

Fleiss JL, Wallenstein S, Rosenfeld R. Adjusting for baseline measurements in the two-period crossover study: a cautionary note. *Control Clin Trials*. 1985 Sep;6(3):192-7.

Hatsukami et al., 2007

D.K. Hatsukami, A.M. Joseph, M. Lesage, J. Jensen, S.E. Murphy, P.R. Pentel, M. Kotlyar, E. Borgida, C. Le, and S.S. Hecht, Developing the science base for reducing tobacco harm, 9(4) *Nicotine & Tobacco research* S537 (Dec. 2007).

Health Canada, 1999

Health Canada. Determination of “Tar”, Nicotine and Carbon, Monoxide in Mainstream Tobacco Smoke. Available at www.hc-sc.gc.ca.

Heatherton et al, 1991

Heatherton TF, Kozlowski LT, Frecker RC, Fagerström KO. The Fagerström test for nicotine dependence: a revision of the Fagerström Tolerance Questionnaire. Br J Addict. 1991;86(9):1119-1127.

Honjo et al. 2010a

Honjo K, Iso H, Tsugane S, Tamakoshi A, Satoh H, Tajima K et al. The effects of smoking and smoking cessation on mortality from cardiovascular disease among Japanese: pooled analysis of three large-scale cohort studies in Japan. Tob Control. 2010; 19 (1): 50-57.

Honjo et al. 2010b

Honjo K, Iso H, Inoue M, Tsugane S; JPHC Study Group. Smoking cessation: predictive factors among middle-aged Japanese. Nicotine Tob Res. 2010; 12 (10): 1050-1054.

ICH E9, 1998

European Medicines Agency. ICH Topic E9: Statistical Principles for Clinical Trials (CPMP/ICH/363/96 – adopted March 1998).

Available at

<http://www.fda.gov/downloads/Drugs/GuidanceComplianceRegulatoryInformation/Guidances/ucm073137.pdf>.

Institute of Medicine, 2012

Institute of Medicine Consensus Report. Scientific Standards for Studies on Modified Risk Tobacco Products. Released: December 14, 2011. Board on Population Health and Public Health Practice. (<http://www.iom.edu/Reports/2011/Scientific-Standards-for-Studies-on-Modified-Risk-Tobacco-Products.aspx>).

International Tobacco Online, 2012

<http://www.internationaltobaccoonline.com/japanese-survey-said-the-percentage-of-smokers-in-japan-is-lower-than-20-p-927.html> (accessed 19 June 2012).

Iso et al. 2005

Iso H, Date C, Yamamoto A, Toyoshima H, Watanabe Y, Kikuchi S et al.; JACC Study Group. Smoking cessation and mortality from cardiovascular disease among Japanese men and women: the JACC Study. Am J Epidemiol. 2005; 161 (2): 170-179.

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

Jacob et al. 2011

Jacob P 3rd, Yu L, Duan M, Ramos L, Yturralde O, Benowitz NL. Determination of the Nicotine Metabolites Cotinine and *Trans*-3'-Hydroxycotinine in Biologic fluids of Smokers and Non-smokers using Liquid Chromatography - Tandem Mass Spectrometry: Biomarkers for Tobacco Smoke Exposure and for Phenotyping Cytochrome P450 2A6 Activity. J Chromatogr B Analyt Technol Biomed Life Sci. 2011 Feb 1; 879 (3-4): 267-76.

Lacy et al. 2007

Lacy CF, Armstrong LL, Goldman MP, Lance LL eds. Drug Information Handbook 15th ed. Hudson, OH; LexiComp Inc. 2007: 1899-1912.

Lingling, 2008

Lingling Han, Merck & Co., Inc., North Wales, PA. Calculating the point estimate and confidence interval of Hodges-Lehmann's median using SAS® software. Paper ST-154, SESUG Proceedings. Available from [http:// analytics.ncsu.edu/sesug/2008/ST-154.pdf](http://analytics.ncsu.edu/sesug/2008/ST-154.pdf) (accessed on 04 Oct 2012)

Matta et al. 2006

Matta SG, Balfour DJ, Benowitz NL, Boyd RT, Buccafusco JJ, Caggiula AR et al. Guidelines on nicotine dose selection for in vivo research. Psychopharmacology (Berl). 2007; 190 (3): 269-319.

Ministry of Health, Labour and Welfare, 1997

Ministerial Ordinance on Good Clinical Practice for Drugs, Ordinance of the Ministry of Health and Welfare No. 28 of March 27, 1997, as last amended by the Ordinance of Ministry of Health, Labour and Welfare No. 161 of December 28, 2012

PMI, 2012a

Philip Morris International. 2013. Internally-modified method based on International Organization for standardization's ISO 4387:2000. In French. PMI_RD_WKI_000530: Détermination du TMP et CO et trappage de l'eau, nicotine, acide pyruvique et des smoke constituents:BaP, phénols, TSNA et amines aromatiques.

PMI, 2012b

Philip Morris International, 2012. In French. Work Instruction PMI_RD_WKI_000527: Guide d'utilisation de l'extracteur in situ, nettoyage des seringues, des pistons, des aiguilles et des pieges (R2540.M109). Version 3.0.

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

PMI, 2013a

Philip Morris International, Neuchatel, Switzerland, 2013. Investigator's Brochure for Tobacco Heating System 2.2.

PMI, 2013b

Philip Morris International. 2013. David Ghosh. Experimental report: Identifying the appropriate technique to collect and extract water, nicotine and glycerin from the TPM to further characterize the tar of the ZURICH product. Version 1.0.

PMI White Paper Docket

Food and Drug Administration (FDA), Division of Dockets Management. Philip Morris's International comments on the scientific evaluation of the modified risk tobacco product applications. Docket no. FDA-2011-N-0443.

Rodgman and Perfetti, 2009

A. Rodgman, and T.A. Perfetti, the chemical components of tobacco and tobacco smoke (Taylor & Francis Group 2009).

Senn, 2002

Senn S. Cross-over Trials in Clinical Research. 2002; John Wiley & Sons. ISBN: 978-0-471-49653-3

Shiffman et al, 2004

Shiffman S, West R, Gilbert D; SRNT Work Group on the Assessment of Craving and Withdrawal in Clinical Trials. Recommendation for the assessment of tobacco craving and withdrawal in smoking cessation trials. Nicotine Tob Res. 2004 Aug;6(4):599-614.

Tiffany et al. 1991

Tiffany ST, Drobes DJ. The development and initial validation of a questionnaire on smoking urges. British Journal of Addiction. 1991; 86: 1467-1476.

WHO, 2002

Evidence Based Recommendations on the Treatment of Tobacco Dependence. June 2001.

Zeller and Hatsukami, 2009

Zeller M, Hatsukami D; Strategic Dialogue on Tobacco Harm Reduction Group. The Strategic Dialogue on Tobacco Harm Reduction: a vision and blueprint for action in the US. Tob Control. 2009; 18.

ZRHX-PK-02, 2012

Philip Morris International, Switzerland. 2012. Clinical study. A Single-center, Open-label, Randomized, Controlled, Crossover Study to Explore the Nicotine Pharmacokinetic Profile and Safety of Tobacco Heating System (THS) 2.1 Compared to Conventional Cigarettes Following Single and ad Libitum Use in Smoking, But Otherwise Healthy Subjects. Available at <http://clinicaltrials.gov/ct2/show/NCT01780688?term=NCT01780688&rank=1>

Appendix 1 Schedule of Events

	Screening	Admission	Wash-out	Single use	Wash-out	Single use	Day of Discharge ^k	Safety follow-up ^l
Study Day	-29 to -2	-1	0	1	2	3	4	4 to 11
Informed consent/subject information sheet	•							
Advice on the risks of smoking and debriefing	•	•					•	
Inclusion/exclusion criteria	•	•						
Enrolment		•						
Randomization			•					
Product test for THS 2.2 Menthol and NRT gum		•						
Product demonstration of THS 2.2 Menthol and NRT gum	•							
Product use				•		•		
Support during periods of reduced smoking/smoking abstinence (as required)			•	•	•	•		
Identification of current mCC brand	•	•						
Smoking history	•	•						
Readiness to abstain from smoking for up to 4 days	•	•						
Willingness to quit smoking in the next 3 months	•							
Demographics ^a , medical history, concomitant diseases	•							
Prior medication ^b /	•	•	•	•	•	•	•	•

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

	Screening	Admission	Wash-out	Single use	Wash-out	Single use	Day of Discharge ^k	Safety follow-up ^l
Study Day	-29 to -2	-1	0	1	2	3	4	4 to 11
Concomitant medication								
Physical examination, body height, weight and related BMI ^c	•	•					•	
Vital signs ^d	•	•	•	•	•	•	•	
ECG	•			•		•		
Spirometry	•	•					•	
Chest X-ray ^e	•							
B/U: Hematology, clinical chemistry, urine analysis	•	•					•	
B: Serology	•							
U: Urine drug screen, urine cotinine screen	•	•						
Alcohol test	•	•						
U: Pregnancy test (all females)	•	•					•	
Collection of used Menthol Tobacco Sticks and mCC butts and NRT gum				•		•		
B: Plasma nicotine ^f				•	•	•	•	
B: COHb ^g				• (5x)		• (5x)		
CO breath test ^h		• (1x)	• (4x)	• (4x)	• (4x)	• (4x)	• (1x)	
B: <i>trans</i> -3'-hydroxycotinine and cotinine (CYP2A6 activity) in plasma		•						
FTND questionnaire	•							
QSU-brief questionnaire ⁱ				•		•		
MCEQ (modified version, only after THS 2.2 Menthol and				•		•		

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

	Screening	Admission	Wash-out	Single use	Wash-out	Single use	Day of Discharge ^k	Safety follow-up ^l
Study Day	-29 to -2	-1	0	1	2	3	4	4 to 11
mCC use								
Cough assessment ^j			•	•	•	•	•	
AE/SAE recording	•	•	•	•	•	•	•	•

Abbreviations: AE = Adverse event; BMI = Body mass index; CO = Carbon monoxide; COHb = Carboxyhemoglobin; CYP2A6 = Cytochrome P450 2A6; ECG = Electrocardiogram; FTND = Fagerström Test for Nicotine Dependence (revised version); mCC = menthol conventional cigarette(s); MCEQ = Modified Cigarette Evaluation Questionnaire; NRT gum = Nicotine replacement therapy gum; QSU-brief = Questionnaire of Smoking Urges brief; SAE = Serious adverse event; THS 2.2 Menthol= Tobacco Heating System 2.2 Menthol

B : Blood sample required.

U : Urine sample required.

a: Sex, date of birth/age

b: Prior medication at Screening and the 4 weeks prior to Screening.

c: Including height (only at Screening), body weight and calculated BMI.

d: Systolic and diastolic blood pressure, pulse rate, respiratory rate.

e: Pre-study chest X-ray (anterior-posterior and left lateral views) to be used, if performed within 6 months prior to Screening.

f: Nicotine blood samples to be taken as follows:

Single use THS 2.2 Menthol and mCC only:

For subjects on THS 2.2 Menthol or on mCC, a total of 16 blood samples will be taken. The first blood sample will be taken within 15 minutes prior to the product use. Thereafter in relation to T₀, blood will be drawn at the following time points: T₁ after 2 min +1 min, T₂ after 4 min +1min, T₃ after 6 min +1 min, T₄ after 8 min +1 min, T₅ after 10 min +1 min, T₆ after 15 min +2 min, T₇ after 30 min +2 min, T₈ after 45 min +2 min, T₉ after 60 min +3 min, T₁₀ after 2 hours +5 min, T₁₁ after 4 hours +5 min, T₁₂ after 6 hours +5 min, T₁₃ after 9 hours +5 min, T₁₄ after 12 hours +5 min, and T₁₅ after 24 hours +5 min.

Single Use on Day 1 and Day 3 for NRT gum only:

For subjects on NRT gum: A total of 16 blood samples will be taken for a 24-hour profile (Day 1 and Day 3). The first blood sample will be taken within 15 min T₀, 15 samples thereafter in relation to T₀: T₁' after 10 min + 1 min, T₂' after 20 minutes + 1 min, T₃' after 25 minutes + 1 min, T₄' after 30 minutes + 1 min, T₅' after 35 minutes + 1 min; T₆' after 40 minutes + 1 min; T₇' after 45 minutes + 1 min, T₈' after 60 minutes + 3 min, T₉' after 2 hours + 5 min, T₁₀' after 3 hours + 5 min, T₁₁' after 4 hours + 5 min, T₁₂' after 6 hours + 5 min, T₁₃' after 9 hours + 5 min, T₁₄' after 12 hours + 5 min, and T₁₅' after 24 hours + 5 min (this sample will be drawn during the day following product use, i.e., wash-out).

g: COHb blood samples to be taken as follows:

Single use: A total of 5 blood samples will be taken. The first sample within 15 minutes prior to T₀ (start of single product use); thereafter in relation to T₀ at 15 min +2 min, 60 min +3 min, 4 hours +5 min and 12 hours +5 min.

h: A CO breath test will be conducted once on Day -1 and Day 4. On Day 0, Day 1, Day 2, Day 3, four breath tests will be done per day. On Day 1 and Day 3, the first test per day will be performed within 15 minutes prior to T₀ (T₀ = start of first product use). The three other tests will be conducted as described in [section 9](#). On the wash-out days (Day 0 and Day 2) the first test will be conducted between 8:00 am-09:30 am. The three other tests will be conducted as described in [section 9](#).

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

i: QSU-brief will be assessed as follows:

Single use: The QSU-brief will be completed by the subject himself/herself at single use study days.

THS 2.2 Menthol and mCC arms: First sample within 15 min prior to T0, 9 assessments thereafter in relation to T0: 15 min, 30 min, 45 min, 60 min, 2 hr, 4 hrs, 6 hrs, 9 hrs, and 12 hrs after T0 with a time window of 10 minutes each.

NRT arm: First sample within 15 min prior to T0, 9 assessments thereafter in relation to T0: 20 min, 30 min, 45 min, 60 min, 2 hr, 4 hr, 6 hrs, 9 hrs, and 12 hrs after T0 with a time window of 10 min each.

j: Visual analogue scale, three Likert scales and one open question.

k: All examinations listed at the Day of Discharge should also be conducted in subjects preliminarily terminating the study.

l: Spontaneous reporting of new AEs/SAEs by the subject and active follow-up of ongoing AEs/SAEs by the site.

Appendix 2 Participating Laboratories

The following Laboratories will be used in the study:

CELERION (USA and Switzerland)

USA: Celerion, 621 Rose Street, 68502, Lincoln, NE, USA

Switzerland: Celerion, Allmendstrasse 32, 8320 Fehraltorf, Switzerland

(b) (4)

A large rectangular grey box redacting information, with the text "(b) (4)" to its left. Below this box are two smaller rectangular grey boxes, also redacting information.

TOKIWA CHEMICAL INDUSTRIES CO.,LTD.

16-22, Kamiikebukuro 4-chome, Toshima-ku, Tokyo, 170-0012, Japan

More details will be found in the study laboratory manuals.

Appendix 3 Investigational Product and Instructions for Use

The product user guide will be provided as a separate document.

Appendix 4 THS Package Label

The package label for the THS Menthol Tobacco Sticks (example):

(b) (4)



Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

The package label for the THS Menthol Tobacco Sticks carton (example):

(b) (4)



Appendix 5 Abnormal Laboratory Values**ABNORMAL LABORATORY VALUES RATING: SERUM CHEMISTRY
PARAMETERS**

Serum Chemistry*	Mild (Grade 1)	Moderate (Grade 2)	Severe (Grade 3)
Sodium – Hyponatremia (mmol/l) ** ⁽¹⁾	<LLN - 130	-	<130 - 120
Sodium – Hypernatremia (mmol/l) ** ⁽¹⁾	>ULN - 150	>150 - 155	>155 - 160; hospitalization indicated
Potassium – Hyperkalaemia (mmol/l)** ⁽¹⁾	>ULN - 5.5	>5.5 - 6.0	>6.0 - 7.0; hospitalization indicated
Potassium – Hypokalaemia (mmol/l) ** ⁽¹⁾	<LLN - 3.0	<LLN - 3.0; symptomatic; intervention indicated	<3.0 - 2.5; hospitalization indicated
Glucose – Hypoglycaemia ** ⁽¹⁾ (mg/dL) (mmol/l)	<LLN – 55; <LLN – 3.0	<55 – 40; <3.0 – 2.2	<40 – 30; <2.2 – 1.7
Blood Urea Nitrogen (BUN) (mg/dL) ⁽²⁾	23 – 26	27 – 31	>31
Glucose – Hyperglycaemia: ** Fasting ⁽¹⁾ (mg/dL) (mmol/l)	>ULN – 160; >ULN – 8.9	>160 -250 > 8.9-13.9	>250 – 500; >13.9 – 27.8 Hospitalization indicated
Creatinine increased** ⁽¹⁾	>1 – 1.5 x baseline; >ULN – 1.5 x ULN	>1.5 – 3.0 x baseline; >1.5 – 3.0 x ULN	>3.0 x baseline; >3.0 – 6.0 x ULN
Albumin – Hypoalbuminaemia** ⁽¹⁾ (g/dL) (g/l)	<LLN – 3; <LLN - 30	<3 – 2; <30 - 20	<2; <20
Total Protein – Hypoproteinaemia ⁽²⁾ (g/dL)	5.5 – 6.0	5.0 – 5.4	<5.0
Alkaline phosphatase increased** ⁽¹⁾	>ULN – 2.5 x ULN	>2.5 – 5.0 x ULN	>5.0 – 20.0 x ULN
ALT / AST increased** ⁽¹⁾	>ULN – 3.0 x ULN	>3.0 – 5.0 x ULN	>5.0 – 20.0 x ULN
Gamma-glutamyl transferase increased ⁽¹⁾	>ULN – 2.5 x ULN	>2.5 – 5.0 x ULN	>5.0 – 20.0 x ULN

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

Blood bilirubin increased ** ⁽¹⁾	>ULN – 1.5 x ULN	>1.5 – 3.0 x ULN	>3.0 – 10.0 ULN
Cholesterol high** ⁽¹⁾ (mg/dL) (mmol/l)	>ULN – 300; >ULN – 7.75	>300-400; >7.75-10.34	>400-500; >10.34-12.92
Triglycerides - Hypertriglyceridemia ⁽¹⁾ (mg/dL) (mmol/l)	150 – 300; 1.71 – 3.42	>300 – 500; >3.42 – 5.70	>500 – 1000; >5.70 – 11.40

Abbreviations: ALT = alanine aminotransferase; AST = aspartate aminotransferase; LLN = lower limit of the normal range; ULN = upper limit of the normal range.

Data Sources:

(1) Common Terminology Criteria for Adverse Events (CTCAE) version 4.03. Toxicity Grading Scale for Healthy Adult and Adolescent Volunteers Enrolled in Preventive Vaccine Clinical Trials, U.S. Department of Health and Human Services, FDA, Center for Biologics Evaluation and Research - Guidance for Industry.

(2) Toxicity Grading Scale for Healthy Adult and Adolescent Volunteers Enrolled in Preventive Vaccine Clinical Trials, U.S. Department of Health and Human Services, FDA, Center for Biologics Evaluation and Research - Guidance for Industry.

* Those parameters that are not listed do not have grading categories in either the CTCAE or the FDA guidance documents and will therefore be reviewed by the Investigator and only reported as an AE if considered to be clinically relevant.

** Where parameters in this table are listed in both the CTCAE and the FDA guidance documents, and each document has different values within each grading category, the grading in CTCAE guidance document predominates.

ABNORMAL LABORATORY VALUES RATING: HEMATOLOGY PARAMETERS

Haematology*	Mild (Grade 1)	Moderate (Grade 2)	Severe (Grade 3)
Haemoglobin (Female) – (g/dL) ⁽¹⁾	11.0 – 12.0	9.5 – 10.9	8.0 – 9.4
change from baseline value – (g/dL) ⁽¹⁾	Any decrease – 1.5	1.6 – 2.0	2.1 – 5.0
Haemoglobin (Male) – (g/dL) ⁽¹⁾	12.5 – 13.5	10.5 – 12.4	8.5 – 10.4
change from baseline value – (g/dL) ⁽¹⁾	Any decrease – 1.5	1.6 – 2.0	2.1 – 5.0
Haemoglobin increase – (g/dL) ⁽²⁾	Increase in >0 – 2 above ULN or above baseline if baseline is above ULN	Increase in >2 – 4 above ULN or above baseline if baseline is above ULN	Increase in >4 above ULN or above baseline if baseline is above ULN
WBC Increase – (cell/mm ³) ⁽¹⁾	10,800 – 15,000	15,001 – 20,000	20,001 – 25,000
WBC Decrease - (cell/mm ³) ^{(2)**}	<LLN – 3000; <LLN – 3.0 x 10 ⁻⁹ /l	<3000 - 2000; <3.0 – 2.0 x 10 ⁻⁹ /l	<2000 - 1000; <2.0 – 1.0 x 10 ⁻⁹ /l
Lymphocytes Increase - (cell/mm ³) ⁽²⁾	-	>4,000 – 20,000	>20,000
Lymphocytes Decrease - (cell/mm ³) ^{(2)**}	<LLN – 800; <LLN – 0.8 x 10 ⁻⁹ /l	<800 - 500; <0.8 – 0.5 x 10 ⁻⁹ /l	<500 - 200; <0.5 – 0.2 x 10 ⁻⁹ /l
Neutrophils Decrease - (cell/mm ³) ^{(2)**}	<LLN – 1500; <LLN – 1.5 x 10 ⁻⁹ /l	<1500 - 1000; <1.5 – 1.0 x 10 ⁻⁹ /l	<1000 - 500; <1.0 – 0.5 x 10 ⁻⁹ /l
Eosinophils - (cell/mm ³) ⁽¹⁾	650 – 1500	1501 - 5000	>5000
Platelets Decrease - (cell/mm ³) ^{(2)**}	<LLN – 75,000; <LLN – 75.0 x 10 ⁻⁹ /l	<75,000 – 50,000; <75.0 – 50.0 x 10 ⁻⁹ /l	<50,000 – 25,000; <50.0 – 25.0 x 10 ⁻⁹ /l

Abbreviations: LLN = Lower limit of the normal range; ULN = Upper limit of the normal range; WBC = White blood cell (count).

Data Source: (1) Toxicity Grading Scale for Healthy Adult and Adolescent Volunteers Enrolled in Preventive Vaccine Clinical Trials, U.S. Department of Health and Human Services, FDA, Center for Biologics Evaluation and Research - Guidance for Industry.

(2) Common Terminology Criteria for Adverse Events (CTCAE) version 4.03.

Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

* Those parameters that are not listed do not have grading categories in either the CTCAE or the FDA guidance documents and will therefore be reviewed by the Investigator and only reported as an AE if considered to be clinically relevant.

** Where parameters in this table are listed in both the CTCAE and the FDA guidance documents, and each document has different values within each grading category, the grading in CTCAE guidance document predominates.

ABNORMAL LABORATORY VALUES RATING: URINALYSIS PARAMETERS

Urine*	Mild (Grade 1)	Moderate (Grade 2)	Severe (Grade 3)
Protein ** ⁽¹⁾	1+ proteinuria; urinary protein <1.0 g/24 hours	2+ proteinuria; urinary protein 1.0-3.4 g/24 hours	Urinary protein ≥3.5 g/24 hours
Glucose ⁽²⁾	Trace	1+	2+
Blood – Haematuria ** ⁽¹⁾	Asymptomatic; clinical or diagnostic observations only; intervention not indicated	Symptomatic; urinary catheter or bladder irrigation indicated; limiting instrumental ADL	Gross haematuria; transfusion, IV medications or hospitalization indicated; elective endoscopic, radiologic or operative intervention indicated; limiting self-care ADL

Abbreviations: ADL = Activities of daily living; IV = Intravenous.

Data Source: (1) Common Terminology Criteria for Adverse Events (CTCAE) version 4.03.

(2) Toxicity Grading Scale for Healthy Adult and Adolescent Volunteers Enrolled in Preventive Vaccine Clinical Trials, U.S. Department of Health and Human Services, FDA, Center for Biologics Evaluation and Research - Guidance for Industry.

* Those parameters that are not listed do not have grading categories in either the CTCAE or the FDA guidance documents and will therefore be reviewed by the Investigator and only reported as an AE if considered to be clinically relevant.

** Where parameters in this table are listed in both the CTCAE and the FDA guidance documents, and each document has different values within each grading category, the grading in CTCAE guidance document predominates.



PMI RESEARCH & DEVELOPMENT

ZRHM-PK-05-JP Clinical Study Protocol Signature Pages

Study Title: A single-center, open-label, randomized, controlled, crossover study to investigate the nicotine pharmacokinetic profile and safety of Tobacco Heating System 2.2 Menthol (THS 2.2 Menthol) following single use in smoking, healthy subjects compared to menthol conventional cigarettes and nicotine gum

Study Number: ZRHM-PK-05-JP

Product Name: Tobacco Heating System 2.2 Menthol (THS 2.2 Menthol)

Principal Investigator and Affiliation: Fumimasa Nobuoka M.D.
Ageo Medical Clinic
3133 Haraichi, Ageo City Saitama 362-0021, Japan

Sponsor: Philip Morris Products S.A.
Quai Jeanrenaud 5
2000 Neuchâtel

Sponsor Signatories: Christelle Haziza, PhD, Manager Clinical Science
Nicola Lama, PhD, Biostatistician
Tamara Koval, MD, Medical Safety Officer

Version: Final 1.0

Date: 21 June 2013

Confidentiality Statement

Confidentiality Statement: Data and information contained in this document are considered to constitute trade secrets and confidential commercial information, and the legal protections provided to such trade secrets and confidential information are hereby claimed under the applicable provisions of applicable law. No part of this document may be publicly disclosed without the written consent of Philip Morris Products S.A.

**PRINCIPAL INVESTIGATOR'S SIGNATURE****PRINCIPAL INVESTIGATOR'S SIGNATURE**

Study Title: A single-center, open-label, randomized, controlled, crossover study to investigate the nicotine pharmacokinetic profile and safety of Tobacco Heating System 2.2 Menthol (THS 2.2 Menthol) following single use in smoking, healthy subjects compared to menthol conventional cigarettes and nicotine gum

Study Number: ZRHM-PK-05-JP

I have read the protocol described above. I agree to comply with all applicable regulations and to conduct the study as described in the protocol.

By signing the protocol, the Investigator agrees to keep all information and documents provided by the Sponsor in strict confidence and to request similar confidentiality from his/her staff. The information provided by the Sponsor to the investigator may not be disclosed to others without direct written authorization from the Sponsor, except to the extent necessary to obtain informed consent from subjects who wish to participate in the study.

Signed: Fumimasa Nobuoka

Date: 26 / Jan / 2013


DR FUMIMASA NOBUOKA M.D.,

**SIGNATURES OF SPONSORS RESPONSIBLE PERSONNEL****SPONSOR SIGNATORIES**


Study Title: A single-center, open-label, randomized, controlled, crossover study to investigate the nicotine pharmacokinetic profile and safety of Tobacco Heating System 2.2 Menthol (THS 2.2 Menthol) following single use in smoking, healthy subjects compared to menthol conventional cigarettes and nicotine gum

Study Number: ZRHR-PK-05-JP


This Clinical Study Protocol was subject to critical review and has been approved by the Sponsor. The following personnel contributed to writing and/or approving this protocol:

Signed: 
Christelle Haziza, PhD, Manager Clinical Science

Date: 21 June 2013

Signed: 
Nicola Lama, PhD, Biostatistician


Date: 25 JUNE 2013

Signed: 
Tamara Koval, MD, Medical Safety Officer


Date: 25 June 2013




16.1.1.2 STUDY PROTOCOL (LOCAL LANGUAGE VERSION)





























































































































































16.1.1.3 PROTOCOL AMENDMENTS

Not applicable.



16.1.1.4 NOTES TO FILE

File to Note

Protocol Title	A single-center, open-label, randomized, controlled, crossover study to investigate the nicotine pharmacokinetic profile and safety of Tobacco Heating System 2.2 Menthol (THS 2.2 Menthol) following single use in smoking, healthy subjects compared to menthol conventional cigarettes and nicotine gum
Protocol No.	ZRHM-PK-05-JP
Item	e-Diary and VAS cough assessment
Date (DD/MMM/YYYY)	22 NOV 2013
<p>Deviation to the protocol was found during the monitoring on 7 Nov 2013. The details are reported in this document.</p> <p>■ Event</p> <p>Among the assessments which are performed with e-Diary, implementation time of 'Assessment of Cough' on Day2 and Day4 deviated from the allowance defined in the protocol.</p> <p>Allowance defined in the protocol is 24 hours minus 5 minutes from T0 of Day1 and Day3</p> <p>■ Subjects</p> <p>123 assessments of 62 randomized subjects</p> <p>Cohort1 – subject number: 0002,0004,0005,0010,0013,0017,0018,0022,0024,0025,0027</p> <p>Cohort2 – subject number: 0031,0035,0036,0039,0040,0043,0045,0049,0050,0051,0052,0054</p> <p>Cohort3 – subject number: 0060,0061,0063,0066,0067,0070,0071,0072,0073,0074,0075,0076,0078,0082,0083,0084</p> <p>Cohort4 – subject number: 0089,0090,0093,0095,0097,0102,0105,0107*,0113,0119</p> <p>Cohort5 – subject number: 0120,0123,0128,0129,0132,0134,0135,0136,0139,0140,0142,0148,0152</p> <p>*Assessment of cough was done until Day2 for subject number 0107 as this subject was discontinued after randomization</p> <p>■ Cause</p> <p>T0 is set for e-Diary on Day1 and Day3 and the cough assessment becomes available only after the alarm ring which occurs after 24 hours+ 5min from T0. At this time point, the implementation time deviated from the allowance; It was concluded that this deviation is not originated from an oversight from the site but from an issue in the EPRO device configuration.</p> <p>■ Corrective action</p> <p>Because this event cannot be avoidable due to the e-Diary programming, no corrective action can be taken at the site.</p>	

Note to File

Protocol Title	A single-center, open-label, randomized, controlled, crossover study to investigate the nicotine pharmacokinetic profile and safety of Tobacco Heating System 2.2 Menthol (THS 2.2 Menthol) following single use in smoking, healthy subjects compared to menthol conventional cigarettes and gum
Protocol No.	ZRHM-PK-05-JP
Site	Ageo Medical Clinic
Item	Source Document Verification (SDV) process for safety laboratory data in EDC
Date (DD/MMM/YYYY)	04/MAR/2014
File	Trial Master file
<p>Description of Issue:</p> <p>Clarification of the Source Document Verification (SDV) process for safety laboratory data in EDC for the above mentioned study</p> <p>The following process will be followed by the CRA for SDV of safety lab data in the EDC system as directed by PMP:</p> <ul style="list-style-type: none"> • The monitor will verify on the paper safety laboratory report the time of collection of the samples, that all parameters have been analyzed as per protocol and that the Principle Investigator (PI) or designee has assessed the results and classified any out of range values for clinical significance and potential adverse events as evidenced by the PI signature and date on the paper safety lab report • When the data is available in EDC, the CRA will SDV only abnormal/out of range results. SDV of abnormal safety lab data in EDC may only be conducted by the CRA once PI assessment has occurred in EDC. If the abnormal lab data is determined to be Not Clinically Significant (NCS) by the PI, the CRA will then mark the entire lab page in the EDC system as SDV. If an abnormal safety lab value is determined to be Clinically Significant (CS) by the PI the CRA will ensure that an adverse event determination has been made by the PI and an Adverse Event eCRF has been completed if applicable. • If the safety lab data results contain all normal values, the CRA will mark the entire eCRF page as SDV in the EDC system • If the safety lab report document indicates an abnormal lab value and PI assessment did not occur, then the CRA will open a query in the eCRF for that data point requesting PI assessment and an adverse event determination. All other normal lab values may be SDV on the eCRF at that time. Once PI assessment has occurred with subsequent completion of an adverse event eCRF if applicable, the CRA may then mark the entire safety lab eCRF as fully SDV in the EDC system. 	

(b) (4)



Signature:

Signature:

Name/Title: Muriel Benzimra/ Clinical Study Manager- Philip Morris Products S.A.

File to Note

Protocol Title	A single-center, open-label, randomized, controlled, crossover study to investigate the nicotine pharmacokinetic profile and safety of Tobacco Heating System 2.2 Menthol (THS 2.2 Menthol) following single use in smoking, healthy subjects compared to menthol conventional cigarettes and nicotine gum
Protocol No.	ZRHM-PK-05-JP
Item	(b) (4) – subject duplicate registration (To be filed in Investigator Study File and Trial Master File)
Date (DD/MMM/YYYY)	27 Dec 2013

The purpose of this file note is to document deviations and corrective/preventive actions for a protocol deviation regarding duplicate registration of subjects. These subjects have been screened twice for the same study under different study numbers. This deviation was detected on 10 Dec 2013 and corrective/preventive actions is for immediate implementation.

1- Description of Deviation:

Double registration of subjects happened to several subjects as listed below :

No.	Subject#	Cohort	ICF date	Status (SF is for "Screen-failure" subject)
1	AGE0008	Cohort 1	1 Aug 2013	SF (not qualified in Spirometer value and cotinine is negative)
	AGE0030	Cohort 2	22 Aug 2013	SF (cotinine is negative)
2	AGE0065	Cohort 3	6 Sep 2013	SF (eligible subjects for cohort 3 were sufficient.)
	AGE0103	Cohort 4	30 Sep 2013	SF (not qualified due to Spirometer value)
3	AGE0087	Cohort 3	6 Sep 2013	SF (eligible subjects for cohort 3 were sufficient.)
	AGE0117	Cohort 4	30 Sep 2013	Discontinue from enrolment (confinement period not met)
4	AGE0110	Cohort 4	30 Sep 2013	SF (not qualified due to Lab test abnormal value)
	AGE0137	Cohort 5	22 Oct 2013	SF (not qualified due to Lab test abnormal value)

2- [Primary cause]

The site misunderstood the protocol which does not allow any re-screening of subject.

The site only considered that "randomized subjects" could not be re-screened. Site only checked for duplication of the information of clinical study participation history (including name and birthdate of subject) when subjects are planned for randomization and not at the time of consent.

In addition, during the site initiation/ investigator meeting, the CRA did not enough highlight this protocol requirement to the site, and it was not clear for the site that the subjects are not allowed to perform any screening test of this study more than once"

The site has received alert emails from (b) (4) (b) (4)) to notify on "possible subject duplication", after system has identified same birth date and gender for 2 registrations. However, site and CRA disregarded these emails –

3- [Corrective/Preventive actions]

These duplications of screening for a same subject occurred at the site will be recorded as a protocol deviation in the appropriate CRF page. The complete list of "duplication" of screening for a subject is reported in the above table

Duplication of screening will be avoided in the future by implementing the following action at four different study conduct stages:

1. At the timepoint of Site initiation visit:

Site training on protocol screening restriction/ requirement will be reinforced.and particularly on the fact that re-screening of a subject already screened in this study is not allowed)

Site training on (b) (4) alert email configuration will be re-inforced

2. At the timepoint of screening subject visit at Informed Consent form signature:

When a subject comes for a screening visit, the site staff at the reception desk verifies systematically that he/she has never taken any ICF/any screening test for this study previously with the registration list of the study subjects existing at the site.

3. At the timepoint of registration to (b) (4)

Site staff should systematically considers alert emails coming from (b) (4) and in case the email content is not clear, site should refer to (b) (4) helpdesk or CRA /sponsor seeking for clarifications.

4. At the timepoint of site monitoring visit

During the eligibility check, the (b) (4) CRA should systematically verify the non-duplication of subject screening versus the medical records/ subjects names.

(b) (4)





NOTE TO FILE (11-AUG-2014)

Clinical Study: ZRHM-PK-05-JP

Study Title:

A single-center, open-label, randomized, controlled, crossover study to investigate the nicotine pharmacokinetic profile and safety of Tobacco Heating System 2.2 Menthol (THS 2.2 Menthol) following single use in smoking, healthy subjects compared to menthol conventional cigarettes and nicotine gum

Investigational Site:

All sites

Sponsor:

Philip Morris Products S.A., Research & Development
Quai Jeanrenaud 5, 2000 Neuchâtel, Switzerland

Author:

Tamara Koval

Subject

Updated Table 4 CYP2A6: Substrates, Inhibitors, and Inducers

Confidentiality Statement

This document is confidential. Disclosure of contents to third parties is not permitted except by written consent of Philip Morris Products S.A.

Purpose The purpose of this note to file is to inform that Table 4 "CYP2A6: Substrates, Inhibitors, and Inducers", which is a part of all Clinical Studies Protocols of P1 is updated in accordance with the Drug Information Handbook, 23d Edition.

Background: At the time of Clinical Protocols Development for P1, Table 4 created, with reference to the Drug Information Handbook 21d Edition. During Clinical Report preparation of the respective studies, the Table was revised based on the latest 23d Edition of 23d Edition of the Drug Information Handbook.

References Drug Information Handbook, 23d Edition

Table 4. CYP 2A6 : Substrates, Inhibitors, Inducers

Inhibitor	Drug Class
Amiodarone	Antiarrhythmic agent
Desipramine	Antidepressant
Isoniazid	Anti-bacterial drug
Ketoconazole	Anti-fungal agent
Letrozole	Anti-estrogen drug
Methoxsalen	Systemic psoralens
Miconazole	Anti-fungal agent
Tranylcypromine	Antidepressant
Inducer	Drug Class
Amobarbital	Barbiturates
Pentobarbital	Barbiturates
Phenobarbital	Barbiturates/anticonvulsant
Rifampin	Antimycobacterial
Secobarbital	Barbiturates
Substrate	Drug Class
Dexmedetomidine	α 2-Adrenoceptor, sedative
Ifosfamide	Anti-cancer, alkylating agent

Sign as applicable by concerned parties:

Name: Tamara Koval _____ Function Medical Safety Officer

Signed: Tamara Koval Date: 11 August 2014



NOTE TO FILE

1 April 2015

Clinical Study Protocol: ZRHM-PK-05-JP

Study Title:

A single-center, open-label, randomized, controlled, crossover study to investigate the nicotine pharmacokinetic profile and safety of Tobacco Heating System 2.2 Menthol (THS 2.2 Menthol) following single use in smoking, healthy subjects compared to menthol conventional cigarettes and nicotine gum

Investigational Site:

Ageo Medical Clinic
3133 Haraichi, Ageo City
Saitama 362-0021, Japan

Version:

1.0

Date:

1 April 2015

Sponsor:

Philip Morris Products S.A., Research & Development
Quai Jeanrenaud 5, 2000 Neuchâtel, Switzerland

Author:

Andrea Donelli

Subject

Change of Medical Safety Officer

Confidentiality Statement

This document is confidential. Disclosure of contents to third parties is not permitted except by written consent of Philip Morris Products S.A.

Purpose

This note to file is created to document a change in the Sponsor's Medical Safety Officer.

Description

On 27 March 2015, Kausar Aamir MD PhD, left the company and therefore her functions for the study as Medical Safety Officer (MSO) was replaced by Patrick Picavet MD.

Impact of the change:

The change of the Sponsor's MSO occurred after study conduct completion, and clinical study report safety review. Changes to the original setup involve the responsibility of acknowledgment and ensuing signature of the final report.


Signatures:

Author:

Andrea Donelli

Function: Clinical Scientist

Company: Philip Morris Products S.A.

Signed: 


Date: 01/APR/2015

Approvers:

Christelle HAZIZA

Function: Manager Clinical Program

Company: Philip Morris Products S.A.

Signed: 

Date: 01 April 2015